



**MINISTRY OF ENVIRONMENT AND FORESTRY
DIRECTORATE GENERAL OF CLIMATE CHANGE**

Published by:
Directorate GHG Inventory and Monitoring Reporting Verification
Directorate General of Climate Change

Manggala Wanabakti Block IV 6th Floor Wing A
Jl. Jend. Gatot Subroto—Jakarta Pusat, DKI Jakarta 10270, Indonesia
Telp./Fax: 021-57903073, Email: tuigrkppi@gmail.com
Website: <http://www.ditjenppi.menlhk.go.id>



EMISSION REDUCTION REPORT

FOR THE INDONESIA-NORWAY PARTNERSHIP



DIRECTORATE GENERAL OF CLIMATE CHANGE
MINISTRY OF ENVIRONMENT AND FORESTRY
REPUBLIC OF INDONESIA
2019



EMISSION REDUCTION REPORT

FOR THE INDONESIA-NORWAY PARTNERSHIP

**DIRECTORATE GENERAL OF CLIMATE CHANGE
MINISTRY OF ENVIRONMENT AND FORESTRY
REPUBLIC OF INDONESIA
2019**

Preface



Indonesia have committed to a new International climate agreement through the ratification of Paris Agreement in 2016. To achieve this commitment, Indonesia and the Kingdom of Norway agree to continue the partnership to further promote the implementation of REDD+ and to protect the remaining natural forests as well as the carbon-rich peatlands in Indonesia, which are part of the Letter of Intent (LoI) between the Government of Indonesia and the Government of Kingdom of Norway that has been signed in 2010.

Since the commencement of the LoI, both countries show strong commitments in addressing issue of climate change mitigation action and in particular supporting the preparedness of REDD+ implementation in Indonesia. Various supports from the government of Norway has been granted to support initiatives in improving capacities and developing systems for implementation of REDD+ and peatland management in Indonesia. Indonesia, as the REDD+ implementing countries, continue to meet the reporting requirement to the UNFCCC. The 1st Forest Reference Emission Level (FREL) has been submitted to and approved by the UNFCCC in 2016. Furthermore, in 2018 Indonesia submitted the Technical Annex of BUR to UNFCCC, which presents the emission reduction results by Indonesia. In paralel, the Indonesia REDD+ Performance report is also upload at Lima REDD+ Hub Website.

As part of the 3rd phase of Indonesia-Norway LoI, both countries have developed the agreed MRV Protocol outlining further mechanism of the result-based payment specifically for the implementation of Indonesia-Norway REDD+ Partnership. The protocol was developed under mutual relationship and common goals to contribute to the international climate agreement through reduction of emissions from tropical deforestation, forest degradation and peatland management.

As the next steps, Indonesia develops the baseline for the result-based payment, as agreed in the MRV Protocol and submits the emission reduction report from the avoided deforestation and forest degradation for the Indonesia-Norway Partnership. We acknowledge the contributions of relevant institutions and team of experts during preparation and development of the RBP baseline and Emission Reduction Report for The Indonesia-Norway Partnership.

Jakarta, May 2019

Dr. Ir. Ruandha Agung Sugardiman, M.Sc
Director General of Climate Change

Table of Contents

| | |
|--|-----|
| PREFACE | i |
| TABLE OF CONTENTS | ii |
| LIST OF TABLES | iii |
| LIST OF FIGURES | iii |
| 1. INTRODUCTION | 1 |
| 2. RESULT BASED PAYMENT BASELINE INDONESIA-NORWAY PARTNERSHIP | 1 |
| 2.1 <i>Definitions Used</i> | 1 |
| 2.1.1 Forest | 2 |
| 2.1.2 Deforestation | 3 |
| 2.1.3 Forest Degradation | 3 |
| 2.1.4 Baseline for Result Based Payment..... | 3 |
| 2.2 <i>Area, Activities and Pools Covered Results</i> | 3 |
| 2.2.1 Area Covered | 3 |
| 2.2.2 Activities Covered | 3 |
| 2.2.3 Pools and Gases | 3 |
| 2.3 <i>Data</i> | 4 |
| 2.3.1 Activity Data | 4 |
| 2.3.2 Emission Factors..... | 4 |
| 2.4 <i>Methodology and Procedures</i> | 4 |
| 2.4.1 Forest Cover Change Analysis | 4 |
| 2.4.2 Reference Period..... | 5 |
| 2.4.3 RBP Baseline Calculation | 5 |
| 2.5 <i>Results of the Construction of RBP Baseline</i> | 5 |
| 2.5.1 Estimates Emission from Deforestation..... | 5 |
| 2.5.2 Estimates Emission from Forest Degradation..... | 6 |
| 2.6 <i>Constructed and Projected RBP Baseline</i> | 6 |
| 3. RESULTS..... | 7 |
| 4. DESCRIPTION OF THE NATIONAL FOREST MONITORING SYSTEM (NFMS) AND THE INSTITUTIONAL ROLES AND RESPONSIBILITIES FOR MRV OF THE RESULTS ... | 8 |
| 5. DEMONSTRATION THAT METHODOLOGIES ARE CONSISTENT WITH RBP BASELINE | 11 |
| 6. NECESSARY INFORMATION THAT ALLOWS FOR THE RECONSTRUCTION OF THE RESULTS | 11 |
| 7. UNCERTAINTY AND BIAS (TEDDY R)..... | 12 |
| 8. PROPOSED RESULT BASED PAYMENT | 13 |
| REFERENCES | 14 |
| ANNEXES | 15 |

List of Tables

| | |
|--|----|
| Table 1. Land cover classes used in the RBP baseline..... | 2 |
| Table 2. Deforestation Emission Factor | 4 |
| Table 3. Forest Degradation Emission Factor | 4 |
| Table 4. Historical (2006/2007 – 2015/2016) and projected (2016/2017 – 2019/2020) annual emission from deforestation and forest degradation (in tCO ₂), calculated using historical data of 2006/2007 – 2015/2016 | 7 |
| Table 5. Emission reduction from avoided deforestation and forest degradation .. | 7 |
| Table 6. Uncertainty calculation for emission from deforestation and forest degradation | 12 |

List of Figures

| | |
|--|----|
| Figure 1. Average annual historical emissions from deforestation expressed in millions tCO ₂ | 5 |
| Figure 2. Average annual historical emissions from forest degradation expressed in millions tCO ₂ | 6 |
| Figure 3. Annual and average annual historical emissions from deforestation and forest degradation (in MtCO ₂) in Indonesia from 2006/2007 to – 2015/2016 and projected emission from 2016/2017 – 2019/2020..... | 6 |
| Figure 4. Annual emissions from deforestation and forest degradation. Pale colours depict historical emissions and green colours depict 2016/2017 emissions..... | 8 |
| Figure 5. General Indonesian Land Cover map workflow..... | 10 |

List of Annexes

| | |
|---|----|
| Annex 1. The calculation of emissions from deforestation and forest degradation | 15 |
| Annex 2. Emissions from peat decomposition | 22 |
| Annex 3. Emissions from peat fire | 26 |

1. Introduction

Government of Indonesia and Government of Kingdom of Norway have been officially agreed to work together in protecting the remaining natural forest in Indonesia, through the “Cooperation on reducing green house gas emissions from deforestation and forest degradation”. The Cooperation was well explained on the Letter of Intent between the Government of Indonesia and the Government of Kingdom of Norway which has been signed on May 26th, 2010. In general, the Government of Kingdom of Norway will provide payment to Government of Indonesia, based on the performance of reducing green house gas emissions from deforestation and forest degradation.

One of the products that should be developed in this Cooperation is Emission Reduction Report that provides information on the reference emission level which will be defined as Result Based Payment (RBP) baseline, and the performance of emission reduction in 2017. This report is developed by referring to MRV protocol for the Indonesia-Norway partnership on climate, forests and peat.

This report is an independent document specifically used for Indonesia-Norway Cooperation on reducing green house gas emissions from deforestation and forest degradation. The RBP baseline in this report is different to Indonesia Forest Reference Emission Level (FREL) that has been submitted to UNFCCC, and passed the technical assessment in 2016. The main different between RBP baseline and FREL is its reference period, where the RBP Baseline used the period of 2006-2016, while FREL used period of 1990-2012.

The Ministry of Environment and Forestry (MoEF) is responsible to develop the emission reduction report and submit it to the Government of Kingdom of Norway. This report will be used as a basis of payments from Government of Kingdom of Norway to the performance or achievement of Government of Indonesia in reducing the green house gas emission from deforestation and forest degradation during 2017-2020.

In addition, based on the document annex for MRV protocol, Indonesia also reported reducing emissions from peatlands. Reports on reducing emissions on peatland include emissions from peat decomposition and emissions from peat fires. Reports for peatlands are presented in Annex 2 and 3 of this report.

2. Result Based Payment Baseline Indonesia-Norway Partnership

2.1 Definitions Used

The definitions used in this document are consistent with those in first FREL. The definitions restated in this document include, among others: definition of forest, deforestation, forest degradation and Baseline for Result Based Payment

2.1.1 Forest

The Government of Indonesia through the Minister of Forestry Decree No. 14/2004 regarding A/R CDM, has set up the definition of forest as “Land with a minimum area of 0.25 hectares that contains trees with canopy cover of at least 30 percent that are capable of reaching a minimum height of 5 meters at maturity” (MoFor, 2004).

Forests used in this document refers to the “working definition”, defined as “a land area of more than 6.25 hectares with trees higher than 5 meters at maturity and a canopy cover of more than 30 percent”. The area span is based on the land-cover maps produced through visual interpretation of satellite images at a scale of 1:50.000 where the minimum area for polygon delineation is 0.25 cm² which equals to 6.25 ha (minimum mapping unit).

Forests used as basis for calculation refer to natural forests, following the classification of Ministry of Environment and Forestry’s land cover map (Table 1) and has been used in the Forest References Emissions Level. The natural forests included six classes, i.e. primary dry land forest, secondary dry land forest, primary swamp forest, secondary swamp forest, primary mangrove forest, and secondary mangrove forest.

Table 1 Land cover classes used in the RBP baseline

| No | Land-cover class | Abbreviation | Category | IPCC |
|-----|---------------------------|--------------|----------------|------------|
| 1. | Primary dryland forest | PF | Natural forest | Forest |
| 2. | Secondary dryland forest | SF | Natural forest | Forest |
| 3. | Primary mangrove forest | PMF | Natural forest | Forest |
| 4. | Secondary mangrove forest | SMF | Natural forest | Forest |
| 5. | Primary swamp forest | PSF | Natural forest | Forest |
| 6. | Secondary swamp forest | SSF | Natural forest | Forest |
| 7. | Plantation forest | TP | Plantation | Forest |
| 8. | Estate crop | EP | Non-forest | Crop land |
| 9. | Pure dry agriculture | AUA | Non-forest | Crop land |
| 10. | Mixed dry agriculture | MxUA | Non-forest | Crop land |
| 11. | Dry shrub | Sr | Non-forest | Grassland |
| 12. | Wet shrub | SSr | Non-forest | Grassland |
| 13. | Savanna and Grasses | Sv | Non-forest | Grassland |
| 14. | Paddy Field | Rc | Non-forest | Crop land |
| 15. | Open swamp | Sw | Non-forest | Wetland |
| 16. | Fish pond/aquaculture | Po | Non-forest | Wetland |
| 17. | Transmigration areas | Tr | Non-forest | Settlement |
| 18. | Settlement areas | Se | Non-forest | Settlement |
| 19. | Port and harbor | Ai | Non-forest | Other land |
| 20. | Mining areas | Mn | Non-forest | Other land |
| 21. | Bare ground | Br | Non-forest | Other land |
| 22. | Open water | WB | Non-forest | Wetland |
| 23. | Clouds and no-data | Ot | Non-forest | No data |

2.1.2 Deforestation

Deforestation is defined as one-time conversion of natural forest cover to other land-cover categories that occurred in the same area. This means that the deforestation occurred in regenerated forests, that previously deforested, are not included in the calculation. This includes conversion of natural forest cover into plantation forest or non-forested lands.

2.1.3 Forest Degradation

Forest degradation is defined as a transition of primary forest classes, which include primary dryland, primary mangrove and primary swamp forests, to secondary forest classes, which reduce the quantity of carbon stocks as a result of human activities. These represent secondary forests that were subject to selective logging or other disturbance events (e.g. fires and encroachment).

2.1.4 Baseline for Result Based Payment

Baseline for result based payment (RBP) is a benchmark for assessing Indonesia's performance in implementing REDD+ under the framework of Norway-Indonesia Partnership. The performance of emission reduction was expressed in tons of carbon dioxide equivalent per year. The technical definition of RBP baseline adopted in this report is a projection of CO₂ gross emissions that is used as a reference to compare against actual emissions at a given point of time in the future. In accordance with MRV protocol document, the RBP baseline will be updated periodically indicatively every 5 years, taking into account any updates of Indonesia's FREL that might be submitted to the UNFCCC. This RBP baseline was developed based on historical forest dynamics and serves as a benchmark for future performance evaluation of REDD+ activities.

2.2 Area, Activities and Pools Covered Results

2.2.1 Area Covered

RBP baseline calculation cover the whole natural forests in Indonesia, which includes dryland, mangrove, and swamp forests from both primary and secondary classes.

2.2.2 Activities Covered

RBP baseline calculation covers the activities related to deforestation and forest degradation. Other REDD+ activities such sustainable management of forest, role of conservation, and enhancement of forest carbon stock were not covered in the calculation.

2.2.3 Pools and Gases

RBP baseline calculation considers aboveground biomass (AGB) as the most significant carbon pool, and reports the greenhouse gas emissions of carbondioxide (CO₂).

2.3 Data

2.3.1 Activity Data

Activity data were generated from the series of land cover maps produced by the Ministry of Environment and Forestry (MoEF), which are the product of the National Forest Monitoring System (NFMS). The maps are accessible via the website (http://webgis.menlhk.go.id:8080/nfms_simontana/). The datasets of 2006, 2009, 2011, 2012, 2013, 2014, 2015, and 2016 land cover maps were used to analyse historical land cover changes, and calculate the emissions estimates.

2.3.2 Emission Factors

RBP baseline calculation uses the emission factors that identical to emission factors used in the 1st Indonesia's FREL. The primary data source used to derive the emission factors were the National Forest Inventory (NFI) - a national program initiated by the Ministry of Forestry in 1989 and supported by the Food and Agriculture Organization of the United Nations (FAO) and the World Bank through the NFI Project. Additionally, research and published data collected from Indonesian sites were used to fill critical data gap currently not available for analysis. Detail emission factor for deforestation and forest degradation see table 2-3.

Table 2 Deforestation Emission Factor

| Forest Classes | Emission Factors of Deforestation (tCO ₂ -e) | | | | | | |
|---------------------------|---|------------|--------|----------|-------|----------|----------|
| | JAWA | KALIMANTAN | MALUKU | NUSABALI | PAPUA | SULAWESI | SUMATERA |
| Primary Dryland Forest | 458,8 | 464,7 | 519,9 | 473,3 | 412,4 | 474,7 | 463,3 |
| Secondary Dryland Forest | 294,1 | 350,7 | 383,1 | 280,6 | 311,2 | 356,2 | 314,3 |
| Primary Mangrove Forest | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 |
| Primay Swamp Forest | 332,5 | 474,0 | 332,5 | 332,5 | 308,4 | 369,8 | 380,9 |
| Secondary Mangrove Forest | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 |
| Secondary Swamp Forest | 274,8 | 294,1 | 274,8 | 274,8 | 251,3 | 221,3 | 261,1 |

Note : If not avaiable data for the emission factor by island, used National Average

Table 3 Forest Degradation Emission Factor

| ForestClasses | Emission Factors of Forest Degradation (tCO ₂ -e) | | | | | | |
|--------------------------|--|------------|--------|----------|-------|----------|----------|
| | JAWA | KALIMANTAN | MALUKU | NUSABALI | PAPUA | SULAWESI | SUMATERA |
| Primary Dryland Forest | 164,7 | 114,0 | 136,8 | 192,7 | 101,3 | 118,5 | 149,0 |
| Secondary Dryland Forest | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 |
| Primary Mangrove Forest | 57,7 | 179,9 | 57,7 | 57,7 | 57,1 | 148,5 | 119,7 |

Note : If not avaiable data for the emission factor by island, used National Average

2.4 Methodology and Procedures

2.4.1 Forest Cover Change Analysis

Annual forest cover change analysis was conducted by overlaying land cover maps of two subsequent periods. Referring to the working definition, deforestation is the change of

natural forests into other classes that occurred one time at any given location across the entire observation period (2006/2007 – 2015/2016).

Forest degradation is the change of primary forests to secondary forests classes in the subsequent year. As elaborated in Margono *et al* (2015), the land cover (LC) data set is a series (T_1 to T_{1+n}) of data, and the degraded forest was generated by comparing the LC of T_n (class of primary forests in the first observation period) to the LC of T_{n+1} (becoming class of secondary forests in the consecutive observation period). Detail information for the calculation process see annex 1.

2.4.2 Reference Period

RBP baseline was calculated based on the period of 2006/2007 – 2015/2016 as the reference period.

2.4.3 RBP Baseline Calculation

RBP baseline calculation was calculated by using the average annual emissions from 2006/2007 to 2015/2016, i.e. from historical emissions from deforestation and forest degradation.

2.5 Results of the Construction of RBP Baseline

2.5.1 Estimates Emission from Deforestation

The averaged annual emission from deforestation in the period 2006/2007 – 2015/2016 is 236.9 MtCO₂ yr⁻¹ (see Figure 1).




Figure 1. Average annual historical emissions from deforestation expressed in millions tCO₂.

2.5.2 Estimates Emission from Forest Degradation

The annual emission from forest degradation in the period 2006/2007 – 2015/2016 is 41.6 MtCO₂ yr⁻¹ (see Figure 2).




Figure 2. Average annual historical emissions from forest degradation expressed in millions tCO₂.

2.6 Constructed and Projected RBP Baseline

The annual total emissions from deforestation and forest degradation in the period of 2006/2007 – 2015/2016 is 278.5 MtCO₂ yr⁻¹ (see Figure 3).




Figure 3. Annual and average annual historical emissions from deforestation and forest degradation (in MtCO₂) in Indonesia from 2006/2007 to – 2015/2016 and projected emission from 2016/2017 – 2019/2020.

Baseline emissions from deforestation and forest degradation are generated based on annual emissions in the period of 2006/2007 to 2015/2016. Detailed annual emissions are shown in the Table 4.

Table 4 Historical (2006/2007 – 2015/2016) and projected (2016/2017 – 2019/2020) annual emission from deforestation and forest degradation (in tCO₂), calculated using historical data of 2006/2007 – 2015/2016

| Year | Deforestation | Forest Degradation | Total annual emission | |
|-----------|---------------|--------------------|-----------------------|------------|
| 2006-2007 | 286.400.629 | 59.226.954 | 345.627.583 | Historical |
| 2007-2008 | 286.400.629 | 59.226.954 | 345.627.583 | |
| 2008-2009 | 286.400.629 | 59.226.954 | 345.627.583 | |
| 2009-2010 | 173.891.040 | 18.511.560 | 192.402.600 | |
| 2010-2011 | 173.891.040 | 18.511.560 | 192.402.600 | |
| 2011-2012 | 248.937.119 | 5.920.802 | 254.857.921 | |
| 2012-2013 | 285.587.006 | 20.395.198 | 305.982.204 | |
| 2013-2014 | 116.066.514 | 9.840.253 | 125.906.767 | |
| 2014-2015 | 232.677.722 | 85.989.932 | 318.667.654 | |
| 2015-2016 | 279.222.082 | 78.664.647 | 357.886.729 | |
| 2016-2017 | 236.947.441 | 41.551.481 | 278.498.922 | Baseline |
| 2017-2018 | 236.947.441 | 41.551.481 | 278.498.922 | |
| 2018-2019 | 236.947.441 | 41.551.481 | 278.498.922 | |
| 2019-2020 | 236.947.441 | 41.551.481 | 278.498.922 | |

3. Results

Emission reductions are calculated through deduction of baseline emission with actual annual emission. In this report, the baseline emissions were generated from the average emission of 2006/2007 – 2015/2016. The baseline for deforestation and forest degradation are 236,947,440 tCO₂.year⁻¹ and 41,551,481 tCO₂.year⁻¹, respectively. While the 2017/2018 actual emission was derived from the current emission from deforestation (228,349,830 tCO₂) and/or forest degradation (42,743,041 tCO₂).

Table 5 Emission reduction from avoided deforestation and forest degradation

| Activity | Emission Reduction (tCO ₂) | Percentage from Baseline (%) |
|--------------------------|--|------------------------------|
| Deforestation | 8,597,611 | 3.63% |
| Forest Degradation | -1,191,560 | -2.87% |
| Total Emission Reduction | 7,406,051 | 2.66% |




Figure 4 Annual emissions from deforestation and forest degradation. Pale colours depict historical emissions and green colours depict 2016/2017 emissions

In 2017, Indonesia has reduced 7,406,051 tCO₂ emission from both avoided deforestation and forest degradation (See Table 5 and figure 4). Avoided emission from 2016/2017 deforestation is 8,597,611 tCO₂ (3.6% from the baseline). However, there is no emission reduction from forest degradation. It is because the actual emission from forest degradation is 2.9% higher than RBP baseline (-1,191,560 tCO₂). It thus deducts the performance from total emission reduction.

4. Description of the National Forest Monitoring System (NFMS) and the institutional roles and responsibilities for MRV of the results

The Ministry of Forestry of Indonesia (MoF) developed forest resource monitoring through National Forest Inventory (NFI) project of Indonesia, established in 1989 (Margono et al., 2016). The NFI project was executed for years under collaboration of the Government of Indonesia (GOI) and Food and Agriculture Organization (FAO). The use of satellite imagery to produce land cover map, which was pre-dominantly Landsat data, was introduced during the periods of NFI. After termination of the NFI project at around 1997/1998, tasks for operationally mapping land cover were transferred into the Forestry Planning Agency/Directorate General (DG) of Forestry Planning of the Ministry of Forestry. The system is now named National Forest Monitoring System (NFMS), which is based on a regular production of land cover map of Indonesia generated in three years interval, and provided in 23 land cover classes including class of cloud cover and no-data. Example of the Indonesia's land cover map is in figure 1, and the National Forest Monitoring System is available online at

http://webgis.menlhk.go.id:8080/nfms_simontana/ for data display, viewing and simple analysis.

Landsat 5 Thematic Mapper (TM) and Landsat 7 Enhanced Thematic Mapper Plus (ETM+) have been used as main data source since early 1990s (Margono et al., 2016). In the tropical region such as Indonesia, clouds and haze are major problems of using optical remotely sensed data, including Landsat. Unluckily, unlike Brazil, Indonesia has no seasonal cloud-free window that offering opportunity to capture clouds-free images [Broich et al 2011 in Margono et al., 2016]. The limited cloud-free image coverage and budget constraint restrict data availability for the system. However since 2008, given the change in Landsat data policy, the United States Geological Survey (USGS) has made Landsat data freely available over the internet [Wulder et al, 2012, Roy et al 2010]. Although most of data are available online at around 2009, the policy significantly gives Indonesia a chance and benefit to increase data available for the system. Total Landsat data to cover the entirety of Indonesia within selected year's interval were approximately 218 scenes.

At the end of 2014, the NFMS established MoU with National Space Agency (LAPAN), particularly in Landsat data provision for ensuring the data sustainability of the NFMS. From that point, LAPAN would automatically provide mosaic of Landsat covering Indonesia (mainly OLI and additional ETM+) on regular basis, which at first would be twice a year. The plan is implemented for 2015 onward.

The 23 land cover classes were generated based on physiognomy or appearance of bio-physical covers that visually distinguished by remote sensing data used: Landsat 30 meter spatial resolution. Detail land-cover category is described in Margono et al., 2016. Although in some extent names of land cover classes mixture to land uses, such as forest plantation or estate crops, object identification over the imagery is purely based on existing appearance, not probable cover or land uses. Several ancillary data sets were used for reference during the process of delineation, to catch as much as valuable information for classification.

Visual classification carried out by digitizing on screen technique using key elements of image/photointerpretation. Under standard GIS software, feature with distinctive existing appearance were visually taken, carefully and manually delineated on the screen to create closed polygons and assigned into designated classes (Figure 4). Recommended maximum scale for classification process is 1:100.000 for using only multispectral bands (e.g. band 5-4-3) and 1:50.000 for using panchromatic band for data registration. A minimum unit polygon is 6.25 hectares or equal to 2.5 cm x 2.5 cm at the maximum zoom screen of 1:50.000 scales or 25 hectares at 1:100.000 scales. Right now, the national land cover map of Indonesia is made available at the scale minimum of 1:250.000.




Figure 5 General Indonesian Land Cover map workflow

QC/QA for the land-cover data was performed using ground-truth points distributed throughout Indonesia. In addition, the assessment of land cover changes uncertainty uses reference data that was generated by using a set of 10,000 of 30x30m grids corresponding to time series of Landsat satellite image pixels (1985-2016). This reference data was selected throughout the country using a simple random sampling technique. The establishment of reference points were also corresponded to other data such as SPOT 6 and 7 satellites from 2013-2016, minimum and maximum values of NDVI, and very high resolution satellite images from Google Earth.

In general, three main problems exist within existing NFMS. Those are (a) presence of persistent clouds cover, (b) inconsistency within the mapping processes, and (c) inability to give near-real time information to match the dynamic recovery of vegetation disturbances. Problems exist due to complex and sometimes conflicting definitions for land cover classification, including forest and deforestation definition. Technically, problems occurred due to lack of adequate data, robust methodologies, and insufficient infrastructure to perform work at a national scale. Adequate data includes timing for data gathering and sufficient pre-processing. Robust methods are linking to rapid needs for supporting carbon monitoring objectives.

The NFMS in a portal is designated to integrate internet ability and forest resource information system in reciprocal (two ways) media information sharing: a step toward the good forest governance, through transparency. As such, information uploaded on NFMS should be maintained, in term of real/near-real time, completeness, and correctness. The current NFMS provides a facility to benefit public participation in updating, correcting, or just commenting the uploaded land cover map. Although currently it might not work as intended, the two ways communication was expected to increase values of correctness, which collected from community in the field as well as from broader users.

5. Demonstration that Methodologies are Consistent with RBP Baseline

This report used a consistent method as in the Chapter 2 on the development of RBP Baseline. This includes consistency in the methodologies used for generating activity data, emission factors, assumptions, definitions, and procedures for estimating CO₂ emissions from deforestation and forest degradation.

Below are specific components used for the emission reduction report that are consistent with the methodologies used for generating the RBP baseline:

- The REDD+ activities included in this report were consistent with the RBP Baseline, i.e. the REDD+ activities with most significant emissions (deforestation and forest degradation).
- The activity data used in this report is the annual land cover map that is generated by the NFMS, which is inline with decision 4/CP 15. This land cover map is produced using the same method as in the RBP Baseline.
- The emission factors used in this report source from the same data used from the RBP Baseline and Indonesia 1st FREL.
- The carbon pools presented in this report were above-ground biomass, maintaining the consistency of the same pools as the RBP Baseline.

6. Necessary Information That Allows For The Reconstruction of The Results

For reconstruction of the results, sources of data needed for the reconstruction of the RBP Baseline and the REDD+ results are provided in the following sites:

1. The data of forest cover, deforestation and degradation that were produced from land cover maps (derived from Landsat imageries) through NFMS for 2006, 2009, 2011, 2012, 2013, 2014, 2015, 2016 & 2017, are accessible online at http://webgis.menlhk.go.id:8080/nfms_simontana/ or <http://geoportal.menlhk.go.id/arcgis/rest/services/KLHK>
2. Other information related the also can access online at <https://geoportal.menlhk.go.id/arcgis/home/>
3. Complete information (spatial data and tables) for the provision of data that allows for the reconstruction of the RBP Baseline and results of the REDD+ can be accessed by request.

Detail information related the reconstruction the RBP baseline and result see annex 1.

7. Uncertainty and Bias

Uncertainty (U) was calculated following the IPCC 2006 Guidelines, volume 1. Chapter 3. If EA is uncertainty from Activity Data and EE is uncertainty from emission factor from activity j, the combined uncertainty (U_j) is calculated using equation :

$$U_j = \sqrt{EA^2 + EE^2} \quad \text{(Equation 1)}$$

Uncertainties from activity data of forest degradation and deforestation were derived from the overall accuracy assessment of land cover map.

A proportion of accuracy contribution (C_j) was calculated from activity j, by involving the uncertainty (U_j), total emissions occurred in the corresponding activities (E_j) and total emission from the corresponding year (E).

$$C_j = (E_j * U_j)^2 / E \quad \text{(Equation 2)}$$

$$TU = \sqrt{\sum C_j} \quad \text{(Equation 3)}$$

Total uncertainty of each year (TU), was derived from a square root of sum C_j.

The uncertainties of emission factor were generated from the standard error of carbon stock values from every forest type/class in each major island/group of islands. The carbon stock was estimated from the NFI plots that have been established in seven major island/group of islands.

The accuracy for the parameter “activity data” (land cover) is 88% (table 6). While the accuracy for the parameter “emission factor” varies from 50-97% depending on island/group of islands and land cover types. The detailed results of the uncertainty analysis for each assessment period are shown in the tables below.

Table 6 Uncertainty calculation for emission from deforestation and forest degradation

| | | | Year | | | | | | | | | | |
|--------------------|--|---------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Component | | Unit | 2006-2007 | 2007-2008 | 2008-2009 | 2009-2010 | 2010-2011 | 2011-2012 | 2012-2013 | 2013-2014 | 2014-2015 | 2015-2016 | 2016-2017 |
| Activity | Deforestation | ton CO2 | 286,400,629 | 286,400,629 | 286,400,629 | 173,891,040 | 173,891,040 | 248,937,119 | 285,587,006 | 116,066,514 | 232,677,722 | 279,222,082 | 228,349,830 |
| | Forest Degradation | ton CO2 | 59,226,954 | 59,226,954 | 59,226,954 | 18,511,560 | 18,511,560 | 5,920,802 | 20,395,198 | 9,840,253 | 85,989,932 | 78,664,647 | 42,743,041 |
| | Total emissions | ton CO2 | 345,627,583 | 345,627,583 | 345,627,583 | 192,402,600 | 192,402,600 | 254,857,921 | 305,982,204 | 125,906,767 | 318,667,654 | 357,886,729 | 271,092,871 |
| Deforestation | AD uncertainty | % | 12 | 12 | 11 | 11 | 11 | 11 | 10 | 10 | 10 | 10 | 10 |
| | EF uncertainty | % | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 |
| | Combined uncertainty | % | 21.63 | 21.63 | 21.14 | 21.14 | 20.99 | 20.88 | 20.79 | 20.74 | 20.67 | 20.57 | 20.57 |
| | Contribution to Variance by Category in Year Base Year | % | 321.35 | 321.35 | 306.77 | 364.93 | 359.75 | 415.92 | 376.65 | 365.49 | 227.88 | 257.48 | 300.13 |
| | Percentage uncertainty in total inventory: | % | 17.9 | 17.9 | 17.5 | 19.1 | 19.0 | 20.4 | 19.4 | 19.1 | 15.1 | 16.0 | 17.3 |
| Forest Degradation | AD uncertainty | % | 12 | 12 | 11 | 11 | 11 | 11 | 10 | 10 | 10 | 10 | 10 |
| | EF uncertainty | % | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 |
| | Combined uncertainty | % | 27.55 | 27.55 | 27.35 | 27.35 | 27.23 | 27.15 | 27.08 | 27.04 | 26.99 | 26.91 | 26.91 |
| | Contribution to Variance by Category in Year Base Year | % | 22.28 | 22.28 | 21.96 | 6.92 | 6.86 | 0.40 | 3.26 | 4.47 | 53.04 | 34.98 | 18.00 |
| | Percentage uncertainty in total inventory: | % | 4.7 | 4.7 | 4.7 | 2.6 | 2.6 | 0.6 | 1.8 | 2.1 | 7.3 | 5.9 | 4.2 |
| Uncertainty | Percentage uncertainty in total inventory: | % | 18.5 | 18.5 | 18.1 | 19.3 | 19.1 | 20.4 | 19.5 | 19.2 | 16.8 | 17.1 | 17.8 |
| | Uncertainty | ton CO2 | 64,070,134 | 64,070,134 | 62,665,139 | 37,102,031 | 36,839,727 | 52,000,651 | 59,639,727 | 24,217,187 | 53,410,429 | 61,204,246 | 48,352,478 |

8. Proposed Result Based Payment

The result based payment baseline for this first reporting period is based on the annual historical average level of each of the following performance indicators: emissions from deforestation and forest degradation. The results based payment baseline for first reporting period is developed using reference period from 2006/2007 – 2015/2016 and valid up to 2019/2020.

Based on MRV Protocol of Norway and Indonesia Partnership, both Parties has agreed terms to treat statistical uncertainty, reversal risk, and possibly other risk factors inclusion of Indonesia's ambition. This treatment term later simplify called set-asides/deductions has been stated in Annex of MRV Protocol that agreed at final meeting between Indonesia and Norway representatives at 7 February 2019. From the reported emission reduction results, the following set-asides/deductions used to determine the maximum number of emission reductions Indonesia can be rewarded for by Norway. The term of set asides/deductions consist of following detail:

- a. From the reported emission reduction results, set-aside/deduction of 20% to reflect the risk of uncertainty in estimates;
- b. In terms of deduction to reflect risk of leakage, Indonesia – Norway agreed to not include this deduction due to the baseline and performance of REDD+ in Indonesia – Norway partnership is counted in national level accounting. Therefor, 0% deduction to reflect risk of leakage is set. The 0% deduction from leakage also consistently used in Indonesia National FREL and REDD+ Performance in 2nd BUR that has been submitted to the UNFCCC as Indonesia approach for implementation REDD+ is in National Level;
- c. In terms of to reflect Indonesia's ambition to reduce national GHG emissions, Indonesia and Norway agreed to deduct 15%.

As systems are developed over time and policies and strategies are put in place to reduce uncertainty risk, risk of leakage, and reflection of Indonesia's ambition, the set aside factor can be reduced. For the first reporting period under the Indonesia – Norway agreement, the total set aside factor of 35% will be applied.

As mentioned on Chapter Results, Indonesia has reduced the emission from deforestation and forest degradation in total amounted to 7,406,051 tCO₂. It comprises of 8,597,611 tCO₂ from reduce deforestation and -1,191,560 tCO₂ from forest degradation. The emission reduction results later deducted 35%. Therefor, Indonesia propose Net results amounted to 4, 813, 933.15 tCO₂ shall be rewarded by Norway for first RBP.

References

- IPCC. 2014. 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands, Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds). Published: IPCC, Switzerland.
- Margono, B.A., Usman, A.B., Budiharto, Sugardiman, R.A. 2016. Indonesia's Forest Resource Monitoring. Indonesian Journal of Geography 48, 7. <https://doi.org/10.22146/ijg.12496>.
- MoEF. 2018. Indonesia Second Biennial Report - Under UNFCCC. http://ditjenppi.menlhk.go.id/reddplus/images/adminppi/dokumen/Indonesia-2nd_BUR_web.pdf.
- MoEF. 2018. MRV Protocol for The Indonesia-Norway Partnership on Climate, Forests and Peat. http://ditjenppi.menlhk.go.id/reddplus/images/adminppi/dokumen/MRV-protocol_final.pdf.
- MoEF. 2016. National Forest Reference Emission Level For Deforestation And Forest Degradation. <http://ditjenppi.menlhk.go.id/reddplus/images/resources/frel/FREL-Submission-by-Indonesia-2016.pdf>.
- MoFor. 2004. *Peraturan Menteri Kehutanan Nomor: P.14/Menhut-II/2004 Tentang Tata Cara Aforestasi dan Reforestasi Dalam Kerangka Mekanisme Pembangunan Bersih*. (p. 1. article 1). Ministry of Forestry. Jakarta.

Annexes

Annex 1. The Calculation of Emission from Deforestation and Forest Degradation

Deforestation and forest degradation emission was calculated using the following equation:

$$GE_{ij} = A_{ij} \times EF_j \quad (1)$$

Where GE_{ij} = CO₂ emissions from deforested or forest degraded area-i at forest change class-j, in tCO₂e. A_{ij} = deforested or forest degradation area-i in forest change class j, in hectares (ha). EF_j = Emission Factor from the loss of carbon stock due to change of forest class-j, owing to deforestation or forest degradation; in tons carbon per ha (tC ha⁻¹). Emission factor from deforestation and forest degradation see table annex 1.1 and table annex 1.2 respectively. Emission from deforestation and forest degradation at period t (GE_t) was estimated using the following equation:

$$GE_t = \sum_{i=1}^N \sum_{j=1}^P GE_{ij} \quad (2)$$

Where, GE_t written in tCO₂, GE_{ij} is emission from deforested or degraded forest area-i in forest class j expressed in tCO₂. N is the number of deforested or degraded forest area unit at period t (from t₀ to t₁), expressed without unit. P is the number of forest classes, which meet natural forest criterion.

Table Annex 1.1. Deforestation Emission Factor

| Forest Classes | Emission Factors of Deforestation (tCO ₂ -e) | | | | | | |
|---------------------------|---|------------|--------|----------|-------|----------|----------|
| | JAWA | KALIMANTAN | MALUKU | NUSABALI | PAPUA | SULAWESI | SUMATERA |
| Primary Dryland Forest | 458,8 | 464,7 | 519,9 | 473,3 | 412,4 | 474,7 | 463,3 |
| Secondary Dryland Forest | 294,1 | 350,7 | 383,1 | 280,6 | 311,2 | 356,2 | 314,3 |
| Primary Mangrove Forest | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 | 455,2 |
| Primay Swamp Forest | 332,5 | 474,0 | 332,5 | 332,5 | 308,4 | 369,8 | 380,9 |
| Secondary Mangrove Forest | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 | 348,0 |
| Secondary Swamp Forest | 274,8 | 294,1 | 274,8 | 274,8 | 251,3 | 221,3 | 261,1 |

*) If not available data for the emission factor by island, used National Average

Table Annex1.2. Forest Degradation Emission Factor

| ForestClasses | Emission Factors of Forest Degradation (tCO ₂ -e) | | | | | | |
|--------------------------|--|------------|--------|----------|-------|----------|----------|
| | JAWA | KALIMANTAN | MALUKU | NUSABALI | PAPUA | SULAWESI | SUMATERA |
| Primary Dryland Forest | 164,7 | 114,0 | 136,8 | 192,7 | 101,3 | 118,5 | 149,0 |
| Secondary Dryland Forest | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 | 107,3 |
| Primary Mangrove Forest | 57,7 | 179,9 | 57,7 | 57,7 | 57,1 | 148,5 | 119,7 |

*) If not available data for the emission factor by island, used National Average

The estimation of emission from deforestation and forest degradation from the loss of above-ground biomass between two years used the land use transition matrix (LUTM). LUTM

derived from the spatial analysis of series of land cover maps, for example series years : 2012 - 2013. Table annex 1.4 provides an example of LUTM transition matrix for the period 2012 - 2013. The emissions from the change of forest change class-j to non-forest classes were calculated using the equation (1). For example, to calculate the emissions from deforestation from primary dryland forest (class code 2001) (GE_{2001}) in tCO₂e, we used the equation (3). Detail class code for the land cover data see table annex 1.3.

$$GE_{2001} = AD * EF \quad (3)$$

Where AD is the change of primary dryland forests (code 2001) to non-forests in the period in hectare; and EF is the emission factor for deforestation of the corresponding class in ton CO₂e/ha (see. Table annex 1.4 and 1.5 presents the sample of the emission matrix from deforestation of all forest classes in 2012-2013).

Table Annex 1.3. Land cover classes used in the Forest Reference Emission Level

| No | Land-cover class | Class Code | Abbreviation |
|-----|---------------------------|------------|--------------|
| 1. | Primary dryland forest | 2001 | PF |
| 2. | Secondary dryland forest | 2002 | SF |
| 3. | Primary mangrove forest | 2004 | PMF |
| 4. | Secondary mangrove forest | 20041 | SMF |
| 5. | Primary swamp forest | 2005 | PSF |
| 6. | Secondary swamp forest | 20051 | SSF |
| 7. | Plantation forest | 2006 | TP |
| 8. | Estate crop | 2010 | EP |
| 9. | Pure dry agriculture | 2009 | AUA |
| 10. | Mixed dry agriculture | 20091 | MxUA |
| 11. | Dry shrub | 2007 | Sr |
| 12. | Wet shrub | 20071 | SSr |
| 13. | Savanna and Grasses | 3000 | Sv |
| 14. | Paddy Field | 20093 | Rc |
| 15. | Open swamp | 50011 | Sw |
| 16. | Fish pond/aquaculture | 20094 | Po |
| 17. | Transmigration areas | 20122 | Tr |
| 18. | Settlement areas | 2012 | Se |
| 19. | Port and harbor | 20121 | Ai |
| 20. | Mining areas | 20141 | Mn |
| 21. | Bare ground | 2014 | Br |
| 22. | Open water | 5001 | WB |
| 23. | Clouds and no-data | 2500 | Ot |

Emissions from the deforestation of other forest classes use similar equation with corresponding emission factors. Therefore the total emission from deforestation of all different forest classes is estimated using the equation (4):

$$GE_t = GE_{2001} + GE_{2002} + GE_{2004} + GE_{2005} + GE_{20041} + GE_{20051} \quad (4)$$

Table Annex 1.4. An example of land use transition matrix of deforestation in the period of 2012-2013 in hectares.

| LC | | LC 2012 (ha) | | | | | | Total |
|---------|-------|--------------|---------|------|--------|--------|---------|---------|
| | | 2001 | 2002 | 2004 | 2005 | .20041 | 20051 | |
| LC 2013 | 2006 | | 12,253 | | 85 | 671 | 39,285 | 52,294 |
| | 2007 | 12,120 | 182,408 | 0 | 242 | | 2,175 | 196,945 |
| | 2010 | 177 | 26,411 | | 858 | 104 | 35,245 | 62,795 |
| | 2012 | | 220 | 46 | 0 | | | 265 |
| | 2014 | 8,690 | 183,816 | 169 | 2,405 | 8,035 | 144,190 | 347,303 |
| | 20071 | 76 | 753 | 741 | 4,254 | 3,768 | 92,907 | 102,499 |
| | 20091 | 188 | 11,553 | | 677 | 47 | 1,725 | 14,190 |
| | 20092 | 9,005 | 121,818 | 0 | 28 | | 1,246 | 132,098 |
| | 20093 | | 283 | | 873 | | | 1,156 |
| | 20094 | | | | 1,672 | | 1,454 | 3,127 |
| | 20121 | | 93 | | | | | 93 |
| | 20122 | | 330 | | | | | 330 |
| | 20141 | 301 | 3,985 | 34 | 268 | 4 | 1,321 | 5,912 |
| | 50011 | | | | | | 241 | 241 |
| Total | | 30,556 | 543,923 | 990 | 11,361 | 12,628 | 319,790 | 919,248 |

Table Annex 1.5. An example of CO₂ emission matrix from deforestation due to loss of above-ground biomass in the period 2012-2013 in tCO₂e.

| LC Classes | | LC 2012 (tCO ₂ e) | | | | | | Total |
|------------|-------|------------------------------|-------------|---------|-----------|-----------|------------|---------------|
| | | 2001 | 2002 | 2004 | 2005 | 20041 | 20051 | |
| LC 2013 | 2006 | - | 4,178,506 | - | 28,287 | 233,445 | 10,794,481 | 15,234,720 |
| | 2007 | 5,560,995 | 62,203,571 | 0 | 80,370 | - | 597,544 | 68,442,479 |
| | 2010 | 81,178 | 9,006,342 | - | 285,338 | 36,153 | 9,684,551 | 19,093,562 |
| | 2012 | - | 74,909 | 20,825 | 0 | - | - | 95,734 |
| | 2014 | 3,986,975 | 62,683,426 | 76,931 | 799,633 | 2,795,735 | 39,619,980 | 109,962,679 |
| | 20071 | 34,773 | 256,747 | 337,179 | 1,414,599 | 1,311,084 | 25,528,741 | 28,883,123 |
| | 20091 | 86,073 | 3,939,799 | - | 225,107 | 16,241 | 474,084 | 4,741,304 |
| | 20092 | 4,131,907 | 41,541,623 | 0 | 9,248 | - | 342,373 | 46,025,152 |
| | 20093 | - | 96,588 | - | 290,157 | - | - | 386,745 |
| | 20094 | - | - | - | 556,083 | - | 399,583 | 955,666 |
| | 20121 | - | 31,768 | - | - | - | - | 31,768 |
| | 20122 | - | 112,529 | - | - | - | - | 112,529 |
| | 20141 | 137,911 | 1,358,785 | 15,547 | 89,137 | 1,255 | 363,087 | 1,965,723 |
| | 50011 | - | - | - | - | - | 66,353 | 66,353 |
| Total | | 14,019,813 | 185,484,592 | 450,481 | 3,777,960 | 4,393,912 | 87,870,778 | 295,997,537 * |

*Note: The total of emission in this calculation is different from the actual emissions in 2013 because this example used the national EF values instead of island-grouping EF values.

For the calculation purposes LUTM as shown in table annex 1.3 summarized by islands, land cover classes and by year period. Detail information for emission from deforestation and forest degradation calculation see table annex 1.5 – 1.8

Table annex 1.5. Activity Data for Deforestation

| Island/Soil/ Land Cover | Deforestation (ha) | | | | | | | |
|---------------------------|--------------------|------------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | 2006-2009 | 2009-2011 | 2011-2012 | 2012-2013 | 2013-2014 | 2014-2015 | 2015-2016 | 2016-2017 |
| SUMATERA | 1.420.549 | 502.062 | 367.706 | 307.579 | 151.461 | 210.212 | 213.258 | 152.982 |
| PEAT | 433.076 | 204.652 | 108.510 | 82.362 | 65.608 | 90.962 | 32.841 | 28.392 |
| Primary Dryland Forest | | | | | | | 613 | 24 |
| Secondary Dryland Forest | 14.920 | 2.708 | 3.691 | 512 | 314 | 986 | | 228 |
| Primary Mangrove Forest | 0 | | | | | 10 | 15 | 1 |
| Secondary Mangrove Forest | 751 | 1.087 | 547 | 50 | 38 | 421 | 654 | 262 |
| Primary Swamp Forest | 37.901 | 10.757 | 5.678 | 5.163 | 1.110 | 9.949 | 10.712 | 703 |
| Secondary Swamp Forest | 379.503 | 190.100 | 98.595 | 76.637 | 64.146 | 79.596 | 20.846 | 27.174 |
| MINERAL | 987.473 | 297.410 | 259.195 | 225.217 | 85.854 | 119.250 | 180.417 | 124.591 |
| Primary Dryland Forest | 8.063 | 7.871 | 7.300 | 7.479 | 8.628 | 2.429 | 22.229 | 14.231 |
| Secondary Dryland Forest | 752.153 | 181.813 | 202.431 | 188.383 | 57.801 | 83.576 | 128.923 | 91.871 |
| Primary Mangrove Forest | 1.043 | 110 | 715 | 145 | 4 | 1.400 | 1.316 | 901 |
| Secondary Mangrove Forest | 24.441 | 2.906 | 5.485 | 1.894 | 1.508 | 4.865 | 11.195 | 6.893 |
| Primary Swamp Forest | 5.001 | 236 | 134 | 492 | 23 | 1.044 | 3.983 | 155 |
| Secondary Swamp Forest | 196.772 | 104.473 | 43.130 | 26.823 | 17.889 | 25.935 | 12.771 | 10.539 |
| KALIMANTAN | 1.021.058 | 458.046 | 292.796 | 494.080 | 154.089 | 348.008 | 423.404 | 315.302 |
| PEAT | 234.606 | 99.684 | 52.164 | 127.764 | 23.146 | 172.957 | 90.678 | 24.439 |
| Primary Dryland Forest | | | | | 16 | | 367 | |
| Secondary Dryland Forest | 5.580 | 1.407 | 2.054 | 3.973 | 184 | 322 | | 415 |
| Primary Mangrove Forest | | 213 | | | | | 0 | |
| Secondary Mangrove Forest | 341 | 19 | 66 | 159 | 20 | 106 | 19 | 210 |
| Primary Swamp Forest | 3.837 | 2.058 | 339 | 4.556 | 503 | 3.446 | 9 | 333 |
| Secondary Swamp Forest | 224.847 | 95.987 | 49.704 | 119.076 | 22.423 | 169.082 | 90.282 | 23.480 |
| MINERAL | 786.452 | 358.362 | 240.632 | 366.317 | 130.944 | 175.051 | 332.726 | 290.863 |
| Primary Dryland Forest | 2.968 | 362 | 6.968 | 11.088 | 1.967 | 1.870 | 5.504 | 1.632 |
| Secondary Dryland Forest | 584.102 | 273.274 | 194.914 | 262.861 | 106.896 | 113.193 | 241.082 | 256.555 |
| Primary Mangrove Forest | 493 | 133 | 164 | 593 | 10 | 116 | 452 | 244 |
| Secondary Mangrove Forest | 22.061 | 3.608 | 8.768 | 5.791 | 3.828 | 5.674 | 11.547 | 7.608 |
| Primary Swamp Forest | 3.237 | 7 | 600 | 122 | 28 | 516 | 577 | 170 |
| Secondary Swamp Forest | 173.591 | 80.977 | 29.219 | 85.863 | 18.214 | 53.682 | 73.563 | 24.654 |
| PAPUA | 115.232 | 31.876 | 43.003 | 23.880 | 22.309 | 81.321 | 17.323 | 51.129 |
| PEAT | 11.987 | 1.729 | 1.039 | 590 | 1.556 | 4.201 | 1.459 | 2.240 |
| Primary Dryland Forest | 48 | 229 | 590 | | 75 | 254 | 98 | 364 |
| Secondary Dryland Forest | 1.848 | 1.359 | 298 | 304 | 473 | 1.490 | 740 | 1.006 |
| Primary Mangrove Forest | 52 | | 37 | 22 | | 18 | 0 | 42 |
| Secondary Mangrove Forest | 212 | 10 | 49 | | | 0 | 0 | 40 |
| Primary Swamp Forest | 4.911 | 105 | 66 | 264 | 642 | 1.309 | 271 | 788 |
| Secondary Swamp Forest | 4.916 | 25 | | | 366 | 1.130 | 350 | |
| MINERAL | 103.246 | 30.147 | 41.964 | 23.290 | 20.753 | 77.121 | 15.863 | 48.889 |
| Primary Dryland Forest | 17.442 | 14.118 | 9.116 | 3.892 | 5.654 | 19.268 | 4.820 | 14.659 |
| Secondary Dryland Forest | 69.499 | 9.952 | 22.597 | 16.312 | 11.243 | 34.274 | 9.438 | 32.327 |
| Primary Mangrove Forest | 49 | 88 | 173 | | 599 | 1.276 | 0 | 134 |
| Secondary Mangrove Forest | 372 | 339 | 238 | 106 | 31 | 165 | 0 | 201 |
| Primary Swamp Forest | 8.403 | 4.974 | 1.532 | 1.931 | 1.129 | 4.859 | 1.422 | 1.513 |
| Secondary Swamp Forest | 7.481 | 677 | 8.308 | 1.049 | 2.097 | 17.279 | 183 | 56 |
| SULAWESI | 140.533 | 74.658 | 19.448 | 46.192 | 16.950 | 56.839 | 91.981 | 77.842 |
| MINERAL | 140.533 | 74.658 | 19.448 | 46.192 | 16.950 | 56.839 | 91.981 | 77.842 |
| Primary Dryland Forest | 4.327 | 18.996 | 1.892 | 6.782 | 1.729 | 6.727 | 17.285 | 6.417 |
| Secondary Dryland Forest | 121.052 | 54.885 | 17.268 | 38.410 | 14.080 | 47.488 | 68.042 | 65.222 |
| Primary Mangrove Forest | 193 | 116 | | 60 | 200 | 60 | 619 | 270 |
| Secondary Mangrove Forest | 3.722 | 556 | 223 | 860 | 708 | 2.221 | 5.131 | 4.247 |
| Primary Swamp Forest | | | | | | 91 | | 3 |
| Secondary Swamp Forest | 11.239 | 105 | 65 | 80 | 233 | 251 | 904 | 1.683 |
| JAWA | 13.244 | 6.100 | 1.294 | 4.349 | 12.976 | 4.495 | 5.015 | 29.863 |
| MINERAL | 13.244 | 6.100 | 1.294 | 4.349 | 12.976 | 4.495 | 5.015 | 29.863 |
| Primary Dryland Forest | 84 | 150 | | | | 81 | | 7 |
| Secondary Dryland Forest | 6.377 | 5.943 | 1.294 | 3.068 | 12.950 | 4.414 | 5.008 | 29.812 |
| Primary Mangrove Forest | | | | | | | 8 | |
| Secondary Mangrove Forest | 6.783 | 7 | | 1.280 | 26 | | 0 | 43 |
| Primary Swamp Forest | | | | | | | | |
| Secondary Swamp Forest | | | | | | | | |
| BALI NUSA | 4.877 | 3.612 | 55.092 | 906 | 1.308 | 18.630 | 30.394 | 9.332 |
| MINERAL | 4.877 | 3.612 | 55.092 | 906 | 1.308 | 18.630 | 30.394 | 9.332 |
| Primary Dryland Forest | 190 | 146 | 1.409 | | 12 | 729 | 3.437 | 623 |
| Secondary Dryland Forest | 4.687 | 3.194 | 52.111 | 864 | 1.288 | 17.512 | 24.493 | 8.664 |
| Primary Mangrove Forest | | 157 | 1.569 | | | 302 | 779 | 10 |
| Secondary Mangrove Forest | | 115 | 3 | 42 | 9 | 87 | 1.684 | 34 |
| Primary Swamp Forest | | | | | | | | |
| Secondary Swamp Forest | | | | | | | | |
| MALUKU | 25.965 | 24.687 | 6.713 | 7.001 | 3.962 | 16.780 | 44.391 | 37.388 |
| MINERAL | 25.965 | 24.687 | 6.713 | 7.001 | 3.962 | 16.780 | 44.391 | 37.388 |
| Primary Dryland Forest | 309 | 1.732 | 10 | 10 | 0 | 599 | 4.476 | 1.033 |
| Secondary Dryland Forest | 25.371 | 21.911 | 6.590 | 6.607 | 3.864 | 15.903 | 36.478 | 33.612 |
| Primary Mangrove Forest | 188 | 1 | 112 | 60 | 75 | 11 | 782 | 650 |
| Secondary Mangrove Forest | 48 | 22 | | 324 | 22 | 225 | 2.522 | 1.449 |
| Primary Swamp Forest | | | | | | 41 | 63 | 23 |
| Secondary Swamp Forest | 50 | 1.021 | | | | | 70 | 622 |
| Grand Total | 2.741.459 | 1.101.040 | 786.052 | 883.986 | 363.056 | 736.285 | 825.766 | 673.838 |
| Annual Rate | 913.820 | 550.520 | 786.052 | 883.986 | 363.056 | 736.285 | 825.766 | 673.838 |

Table annex 1.6. Emission from Deforestation

| Island/Soil/ Land Cover | Emission of Deforestation (t CO2e/Periode) | | | | | | | |
|---------------------------|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| | 2006-2009 | 2009-2011 | 2011-2012 | 2012-2013 | 2013-2014 | 2014-2015 | 2015-2016 | 2016-2017 |
| SUMATERA | 420.883.277 | 144.191.690 | 109.810.804 | 92.745.350 | 44.656.423 | 61.929.177 | 70.205.419 | 48.624.920 |
| PEAT | 118.492.775 | 54.970.891 | 29.260.741 | 22.158.419 | 17.286.314 | 25.036.735 | 10.042.414 | 7.538.507 |
| Primary Dryland Forest | - | - | - | - | - | - | 284.071 | 11.246 |
| Secondary Dryland Forest | 4.689.089 | 851.204 | 1.159.915 | 160.932 | 98.741 | 309.837 | - | 71.661 |
| Primary Mangrove Forest | 39 | - | - | - | - | 4.745 | 7.015 | 357 |
| Secondary Mangrove Forest | 261.438 | 378.070 | 190.166 | 17.396 | 13.237 | 146.388 | 227.560 | 91.095 |
| Primary Swamp Forest | 14.434.966 | 4.097.021 | 2.162.697 | 1.966.415 | 422.655 | 3.789.269 | 4.079.744 | 267.610 |
| Secondary Swamp Forest | 99.107.243 | 49.644.596 | 25.747.962 | 20.013.676 | 16.751.682 | 20.786.496 | 5.444.024 | 7.096.538 |
| MINERAL | 302.390.501 | 89.220.800 | 80.550.063 | 70.586.930 | 27.370.109 | 36.892.441 | 60.163.005 | 41.086.413 |
| Primary Dryland Forest | 3.735.588 | 3.646.567 | 3.382.225 | 3.465.079 | 3.997.422 | 1.125.595 | 10.298.983 | 6.593.370 |
| Secondary Dryland Forest | 236.384.152 | 57.139.657 | 63.619.388 | 59.204.427 | 18.165.666 | 26.265.891 | 40.517.552 | 28.873.039 |
| Primary Mangrove Forest | 474.712 | 50.144 | 325.273 | 66.050 | 1.841 | 637.533 | 598.976 | 410.272 |
| Secondary Mangrove Forest | 8.504.403 | 1.011.233 | 1.908.662 | 659.179 | 524.669 | 1.692.790 | 3.895.212 | 2.398.385 |
| Primary Swamp Forest | 1.904.686 | 90.033 | 51.183 | 187.292 | 8.709 | 397.743 | 1.517.033 | 59.002 |
| Secondary Swamp Forest | 51.386.961 | 27.283.165 | 11.263.332 | 7.004.902 | 4.671.802 | 6.772.889 | 3.335.249 | 2.752.345 |
| KALIMANTAN | 336.715.544 | 150.934.121 | 99.113.621 | 163.552.430 | 52.018.018 | 110.131.543 | 139.964.049 | 108.096.594 |
| PEAT | 70.020.864 | 29.801.789 | 15.521.875 | 38.627.665 | 6.911.763 | 51.509.673 | 26.733.427 | 7.281.992 |
| Primary Dryland Forest | - | - | - | - | 7.352 | - | 170.762 | - |
| Secondary Dryland Forest | 1.956.875 | 493.569 | 720.310 | 1.393.281 | 64.510 | 113.016 | - | 145.684 |
| Primary Mangrove Forest | - | 96.793 | - | - | - | - | 0 | - |
| Secondary Mangrove Forest | 118.808 | 6.585 | 22.874 | 55.250 | 6.916 | 36.935 | 6.677 | 73.165 |
| Primary Swamp Forest | 1.818.590 | 975.570 | 160.870 | 2.159.536 | 238.517 | 1.633.405 | 4.475 | 157.742 |
| Secondary Swamp Forest | 66.126.590 | 28.229.273 | 14.617.820 | 35.019.598 | 6.594.467 | 49.726.316 | 26.551.513 | 6.905.401 |
| MINERAL | 266.694.680 | 121.132.332 | 83.591.746 | 124.924.765 | 45.106.256 | 58.621.870 | 113.230.621 | 100.814.602 |
| Primary Dryland Forest | 1.379.197 | 168.346 | 3.237.846 | 5.152.481 | 914.135 | 868.794 | 2.557.852 | 758.506 |
| Secondary Dryland Forest | 204.828.256 | 95.829.702 | 68.350.879 | 92.177.862 | 37.485.591 | 39.693.652 | 84.540.815 | 89.966.547 |
| Primary Mangrove Forest | 224.391 | 60.737 | 74.553 | 269.725 | 4.741 | 52.854 | 205.919 | 111.072 |
| Secondary Mangrove Forest | 7.676.186 | 1.255.286 | 3.050.840 | 2.014.893 | 1.331.805 | 1.974.440 | 4.017.905 | 2.647.275 |
| Primary Swamp Forest | 1.534.207 | 3.290 | 284.575 | 57.889 | 13.289 | 244.571 | 273.593 | 80.496 |
| Secondary Swamp Forest | 51.052.443 | 23.814.970 | 8.593.053 | 25.251.915 | 5.356.695 | 15.787.558 | 21.634.538 | 7.250.706 |
| PAPUA | 36.885.199 | 11.341.006 | 13.903.308 | 7.763.045 | 7.457.005 | 26.355.263 | 5.851.738 | 17.455.469 |
| PEAT | 3.442.148 | 559.777 | 390.045 | 186.171 | 467.894 | 1.264.206 | 442.373 | 739.077 |
| Primary Dryland Forest | 19.763 | 94.536 | 243.141 | - | 30.771 | 104.686 | 40.496 | 149.924 |
| Secondary Dryland Forest | 575.023 | 423.014 | 92.586 | 94.537 | 147.045 | 463.563 | 230.388 | 313.153 |
| Primary Mangrove Forest | 23.507 | - | 16.751 | 10.182 | - | 8.241 | 0 | 18.958 |
| Secondary Mangrove Forest | 73.782 | 3.528 | 17.209 | - | - | 1 | 0 | 14.060 |
| Primary Swamp Forest | 1.514.477 | 32.378 | 20.357 | 81.452 | 198.043 | 403.850 | 83.632 | 242.983 |
| Secondary Swamp Forest | 1.235.595 | 6.322 | - | - | 92.036 | 283.865 | 87.857 | - |
| MINERAL | 33.443.051 | 10.781.229 | 13.513.264 | 7.576.874 | 6.989.110 | 25.091.057 | 5.409.366 | 16.716.392 |
| Primary Dryland Forest | 7.193.579 | 5.822.406 | 3.759.747 | 1.605.167 | 2.331.955 | 7.946.769 | 1.987.875 | 6.045.666 |
| Secondary Dryland Forest | 21.626.292 | 3.096.742 | 7.031.483 | 5.075.787 | 3.498.439 | 10.665.161 | 2.936.697 | 10.059.389 |
| Primary Mangrove Forest | 22.238 | 39.938 | 78.804 | - | 272.725 | 580.833 | 49 | 60.953 |
| Secondary Mangrove Forest | 129.426 | 118.127 | 82.826 | 36.720 | 10.638 | 57.324 | 0 | 69.847 |
| Primary Swamp Forest | 2.591.521 | 1.533.952 | 472.547 | 595.489 | 348.343 | 1.498.536 | 438.648 | 466.502 |
| Secondary Swamp Forest | 1.879.994 | 170.062 | 2.087.856 | 263.710 | 527.011 | 4.342.434 | 46.096 | 14.034 |
| SULAWESI | - | - | - | - | - | - | - | - |
| MINERAL | 49.042.074 | 28.836.263 | 7.141.031 | 17.244.853 | 6.224.841 | 20.997.728 | 34.708.041 | 28.251.896 |
| Primary Dryland Forest | 2.054.188 | 9.017.254 | 898.220 | 3.219.442 | 820.936 | 3.193.482 | 8.204.968 | 3.046.245 |
| Secondary Dryland Forest | 43.117.649 | 19.549.660 | 6.150.751 | 13.681.195 | 5.015.265 | 16.914.895 | 24.235.884 | 23.231.447 |
| Primary Mangrove Forest | 88.027 | 52.621 | - | 27.198 | 90.942 | 27.261 | 281.619 | 122.905 |
| Secondary Mangrove Forest | 1.294.928 | 193.456 | 77.758 | 299.280 | 246.201 | 772.943 | 1.785.439 | 1.477.678 |
| Primary Swamp Forest | - | - | - | - | - | 33.663 | - | 1.073 |
| Secondary Swamp Forest | 2.487.282 | 23.271 | 14.302 | 17.738 | 51.498 | 55.485 | 200.130 | 372.547 |
| JAWA | - | - | - | - | - | - | - | - |
| MINERAL | 4.274.374 | 1.819.032 | 380.515 | 1.347.901 | 3.817.679 | 1.335.358 | 1.476.122 | 8.785.955 |
| Primary Dryland Forest | 38.733 | 68.813 | - | - | - | 37.158 | - | 3.440 |
| Secondary Dryland Forest | 1.875.450 | 1.747.742 | 380.515 | 902.375 | 3.808.571 | 1.298.199 | 1.472.697 | 8.767.708 |
| Primary Mangrove Forest | - | - | - | - | - | - | 3.416 | - |
| Secondary Mangrove Forest | 2.360.191 | 2.477 | - | 445.526 | 9.108 | - | 9 | 14.806 |
| Primary Swamp Forest | - | - | - | - | - | - | - | - |
| Secondary Swamp Forest | - | - | - | - | - | - | - | - |
| BALI NUSA | - | - | - | - | - | - | - | - |
| MINERAL | 1.405.503 | 1.076.968 | 16.006.653 | 257.243 | 370.065 | 5.427.472 | 9.441.304 | 2.742.926 |
| Primary Dryland Forest | 90.165 | 68.980 | 666.880 | - | 5.528 | 345.190 | 1.626.904 | 294.696 |
| Secondary Dryland Forest | 1.315.338 | 896.307 | 14.624.485 | 242.544 | 361.353 | 4.914.636 | 6.873.880 | 2.431.600 |
| Primary Mangrove Forest | - | 71.684 | 714.153 | - | - | 137.275 | 354.420 | 4.641 |
| Secondary Mangrove Forest | - | 39.996 | 1.134 | 14.699 | 3.184 | 30.371 | 586.100 | 11.988 |
| Primary Swamp Forest | - | - | - | - | - | - | - | - |
| Secondary Swamp Forest | - | - | - | - | - | - | - | - |
| MALUKU | - | - | - | - | - | - | - | - |
| MINERAL | 9.995.918 | 9.583.000 | 2.581.187 | 2.676.184 | 1.522.482 | 6.501.181 | 17.575.409 | 14.392.072 |
| Primary Dryland Forest | 160.574 | 900.288 | 5.386 | 5.101 | 0 | 311.499 | 2.326.890 | 536.952 |
| Secondary Dryland Forest | 9.719.546 | 8.394.018 | 2.524.810 | 2.531.081 | 1.480.491 | 6.092.518 | 13.974.744 | 12.876.639 |
| Primary Mangrove Forest | 85.626 | 465 | 50.990 | 27.131 | 34.368 | 5.072 | 356.039 | 295.948 |
| Secondary Mangrove Forest | 16.530 | 7.626 | - | 112.872 | 7.623 | 78.389 | 877.473 | 504.233 |
| Primary Swamp Forest | - | - | - | - | - | 13.703 | 20.914 | 7.487 |
| Secondary Swamp Forest | 13.642 | 280.602 | - | - | - | - | 19.349 | 170.814 |
| Grand Total | 859.201.888 | 347.782.079 | 248.937.119 | 285.587.006 | 116.066.514 | 232.677.722 | 279.222.082 | 228.349.830 |
| Annual Rate | 286.400.629 | 173.891.040 | 248.937.119 | 285.587.006 | 116.066.514 | 232.677.722 | 279.222.082 | 228.349.830 |

Table annex 1.7 Activity Data for Forest Degradation

| Island/Soil/ Land Cover | Forest Degradation (ha) | | | | | | | |
|-------------------------|-------------------------|----------------|---------------|----------------|---------------|----------------|----------------|----------------|
| | 2006-2009 | 2009-2011 | 2011-2012 | 2012-2013 | 2013-2014 | 2014-2015 | 2015-2016 | 2016-2017 |
| SUMATERA | 70.409 | 45.463 | 2.346 | 16.598 | 1.166 | 39.162 | 228.412 | 94.172 |
| PEAT | 33.571 | 15.421 | 2.228 | 3.943 | 248 | 529 | 24.439 | 96 |
| Primary Dryland Forest | | | | | 248 | | | 5 |
| Primary Mangrove Forest | 258 | | | | | 381 | 81 | 20 |
| Primary Swamp Forest | 33.313 | 15.421 | 2.228 | 3.943 | | 149 | 24.358 | 71 |
| MINERAL | 36.838 | 30.042 | 118 | 12.654 | 917 | 38.633 | 203.973 | 94.076 |
| Primary Dryland Forest | 3.595 | 24.480 | 26 | 1.230 | 774 | 26.598 | 185.991 | 91.761 |
| Primary Mangrove Forest | 28.134 | 2.939 | | 600 | | 11.494 | 3.552 | 2.044 |
| Primary Swamp Forest | 5.109 | 2.624 | 93 | 10.824 | 144 | 541 | 14.431 | 271 |
| KALIMANTAN | 70.608 | 18.019 | 10.210 | 7.751 | 37.644 | 163.876 | 74.153 | 88.304 |
| PEAT | 740 | 166 | 10.210 | 434 | 1.209 | 9.064 | 1.569 | 3.223 |
| Primary Dryland Forest | | | 10.210 | | | 37 | | |
| Primary Mangrove Forest | | | | | 75 | | | 1.330 |
| Primary Swamp Forest | 740 | 166 | | 434 | 1.135 | 9.027 | 1.569 | 1.893 |
| MINERAL | 69.868 | 17.853 | | 7.317 | 36.434 | 154.811 | 72.584 | 85.081 |
| Primary Dryland Forest | 67.975 | 17.713 | | 6.157 | 35.782 | 145.535 | 70.604 | 81.102 |
| Primary Mangrove Forest | 1.887 | | | 284 | 442 | 238 | 1.288 | 3.650 |
| Primary Swamp Forest | 7 | 140 | | 875 | 209 | 9.038 | 691 | 329 |
| PAPUA | 992.217 | 62.177 | 6.165 | 168.199 | 52.894 | 263.144 | 162.406 | 74.612 |
| PEAT | 47.726 | 5.941 | 710 | 14.287 | 4.264 | 8.741 | 5.965 | 2.506 |
| Primary Dryland Forest | 14.533 | 535 | | 4.573 | 330 | 8.111 | 2.199 | 1.793 |
| Primary Mangrove Forest | 3.205 | 255 | | 3.887 | 4 | 325 | 1.084 | 7 |
| Primary Swamp Forest | 29.988 | 5.151 | 710 | 5.828 | 3.930 | 306 | 2.682 | 706 |
| MINERAL | 944.491 | 56.236 | 5.455 | 153.912 | 48.630 | 254.402 | 156.442 | 72.106 |
| Primary Dryland Forest | 817.699 | 37.989 | 1.009 | 138.898 | 29.573 | 249.465 | 135.226 | 64.982 |
| Primary Mangrove Forest | 5.547 | 53 | | 2.642 | 2.769 | 568 | 2.354 | 363 |
| Primary Swamp Forest | 121.244 | 18.194 | 4.445 | 12.372 | 16.288 | 4.369 | 18.862 | 6.761 |
| SULAWESI | 97.610 | 186.799 | 10.462 | 9.113 | 4.706 | 112.521 | 63.622 | 21.274 |
| MINERAL | 97.610 | 186.799 | 10.462 | 9.113 | 4.706 | 112.521 | 63.622 | 21.274 |
| Primary Dryland Forest | 95.666 | 186.707 | 10.462 | 9.113 | 3.250 | 111.322 | 63.334 | 20.734 |
| Primary Mangrove Forest | 1.944 | 92 | | | 1.457 | 850 | 282 | 540 |
| Primary Swamp Forest | | | | | | 349 | 7 | |
| JAWA | 267.460 | | | | 43 | 1.021 | 242 | 308 |
| MINERAL | 267.460 | | | | 43 | 1.021 | 242 | 308 |
| Primary Dryland Forest | 266.518 | | | | 43 | 1.021 | 107 | 31 |
| Primary Mangrove Forest | 942 | | | | | | 87 | 277 |
| Primary Swamp Forest | | | | | | | 48 | |
| BALI NUSA | 59.491 | 2.107 | 15.010 | 255 | 1.158 | 75.660 | 46.856 | 45.813 |
| MINERAL | 59.491 | 2.107 | 15.010 | 255 | 1.158 | 75.660 | 46.856 | 45.813 |
| Primary Dryland Forest | 59.457 | 2.107 | 14.387 | 255 | 1.135 | 74.541 | 44.739 | 44.248 |
| Primary Mangrove Forest | 33 | | 624 | | 23 | 1.119 | 2.117 | 1.565 |
| Primary Swamp Forest | | | | | | | | |
| MALUKU | 5.266 | 7.460 | | 153 | 405 | 48.015 | 39.764 | 1.468 |
| MINERAL | 5.266 | 7.460 | | 153 | 405 | 48.015 | 39.764 | 1.468 |
| Primary Dryland Forest | 56 | 7.375 | | 0 | 41 | 45.665 | 38.719 | 562 |
| Primary Mangrove Forest | 5.210 | 85 | | 153 | 364 | 1.628 | 928 | 716 |
| Primary Swamp Forest | | | | | | 722 | 117 | 189 |
| Grand Total | 1.563.061 | 322.024 | 44.193 | 202.070 | 98.015 | 703.398 | 615.456 | 325.951 |
| Annual Rate | 521.020 | 161.012 | 44.193 | 202.070 | 98.015 | 703.398 | 615.456 | 325.951 |

Table annex 1.8. Emission from Forest Degradation

| Island/Soil/ Land Cover | Emission of Forest Degradatin (t CO2e/Periode) | | | | | | | |
|-------------------------|--|-------------------|------------------|-------------------|------------------|-------------------|-------------------|-------------------|
| | 2006-2009 | 2009-2011 | 2011-2012 | 2012-2013 | 2013-2014 | 2014-2015 | 2015-2016 | 2016-2017 |
| SUMATERA | 8.180.804 | 6.123.553 | 281.552 | 2.015.499 | 169.487 | 5.320.279 | 32.751.495 | 13.938.310 |
| PEAT | 4.015.537 | 1.845.991 | 266.654 | 472.061 | 37.001 | 58.608 | 2.924.541 | 11.362 |
| Primary Dryland Fc | - | - | - | - | 37.001 | - | - | 690 |
| Primary Mangrove | 27.715 | - | - | - | - | 40.826 | 8.682 | 2.172 |
| Primary Swamp Fo | 3.987.823 | 1.845.991 | 266.654 | 472.061 | - | 17.781 | 2.915.858 | 8.500 |
| MINERAL | 4.165.267 | 4.277.562 | 14.898 | 1.543.438 | 132.487 | 5.261.672 | 29.826.955 | 13.926.948 |
| Primary Dryland Fc | 535.760 | 3.648.214 | 3.819 | 183.317 | 115.301 | 3.963.940 | 27.718.485 | 13.675.261 |
| Primary Mangrove | 3.017.890 | 315.264 | - | 64.405 | - | 1.232.978 | 381.020 | 219.252 |
| Primary Swamp Fo | 611.617 | 314.083 | 11.078 | 1.295.717 | 17.186 | 64.753 | 1.727.450 | 32.435 |
| KALIMANTAN | 8.086.937 | 2.074.535 | 1.164.151 | 968.117 | 4.377.040 | 19.873.083 | 8.594.843 | 10.180.890 |
| PEAT | 133.138 | 29.841 | 1.164.151 | 78.140 | 212.145 | 1.628.240 | 282.251 | 483.215 |
| Primary Dryland Fc | - | - | 1.164.151 | - | - | 4.262 | - | - |
| Primary Mangrove | - | - | - | - | 8.037 | - | - | 142.674 |
| Primary Swamp Fo | 133.138 | 29.841 | - | 78.140 | 204.109 | 1.623.978 | 282.251 | 340.541 |
| MINERAL | 7.953.798 | 2.044.694 | - | 889.977 | 4.164.895 | 18.244.844 | 8.312.591 | 9.697.675 |
| Primary Dryland Fc | 7.750.208 | 2.019.584 | - | 701.985 | 4.079.749 | 16.593.359 | 8.050.008 | 9.246.863 |
| Primary Mangrove | 202.364 | - | - | 30.501 | 47.466 | 25.571 | 138.189 | 391.559 |
| Primary Swamp Fo | 1.227 | 25.110 | - | 157.491 | 37.680 | 1.625.913 | 124.395 | 59.253 |
| PAPUA | 93.838.185 | 5.266.497 | 396.533 | 16.266.071 | 4.479.502 | 26.442.652 | 15.513.275 | 7.227.115 |
| PEAT | 3.527.439 | 375.587 | 40.539 | 1.212.667 | 258.234 | 873.525 | 492.046 | 222.642 |
| Primary Dryland Fc | 1.471.483 | 54.155 | - | 463.009 | 33.453 | 821.203 | 222.620 | 181.574 |
| Primary Mangrove | 343.802 | 27.364 | - | 416.927 | 383 | 34.847 | 116.311 | 766 |
| Primary Swamp Fo | 1.712.154 | 294.067 | 40.539 | 332.731 | 224.398 | 17.476 | 153.115 | 40.302 |
| MINERAL | 90.310.746 | 4.890.910 | 355.994 | 15.053.404 | 4.221.268 | 25.569.126 | 15.021.230 | 7.004.473 |
| Primary Dryland Fc | 82.793.340 | 3.846.429 | 102.203 | 14.063.626 | 2.994.316 | 25.258.734 | 13.691.810 | 6.579.534 |
| Primary Mangrove | 595.056 | 5.716 | - | 283.386 | 297.029 | 60.953 | 252.498 | 38.902 |
| Primary Swamp Fo | 6.922.350 | 1.038.765 | 253.791 | 706.391 | 929.922 | 249.439 | 1.076.921 | 386.037 |
| SULAWESI | - | - | - | - | - | - | - | - |
| MINERAL | 11.545.054 | 22.134.776 | 1.239.767 | 1.079.892 | 541.357 | 13.282.864 | 7.535.324 | 2.514.874 |
| Primary Dryland Fc | 11.336.537 | 22.124.946 | 1.239.767 | 1.079.892 | 385.094 | 13.191.710 | 7.505.074 | 2.456.982 |
| Primary Mangrove | 208.517 | 9.830 | - | - | 156.263 | 91.154 | 30.249 | 57.892 |
| Primary Swamp Fo | - | - | - | - | - | - | - | - |
| JAWA | - | - | - | - | - | - | - | - |
| MINERAL | 44.004.030 | - | - | - | 7.155 | 168.132 | 26.901 | 34.876 |
| Primary Dryland Fc | 43.903.005 | - | - | - | 7.155 | 168.132 | 17.558 | 5.165 |
| Primary Mangrove | 101.025 | - | - | - | - | - | 9.343 | 29.710 |
| Primary Swamp Fo | - | - | - | - | - | - | - | - |
| BALI NUSA | - | - | - | - | - | - | - | - |
| MINERAL | 11.459.326 | 405.877 | 2.838.799 | 49.185 | 221.093 | 14.481.974 | 8.847.070 | 8.693.215 |
| Primary Dryland Fc | 11.455.750 | 405.877 | 2.771.894 | 49.185 | 218.606 | 14.361.912 | 8.619.999 | 8.525.321 |
| Primary Mangrove | 3.576 | - | 66.904 | - | 2.488 | 120.062 | 227.071 | 167.894 |
| Primary Swamp Fo | - | - | - | - | - | - | - | - |
| MALUKU | - | - | - | - | - | - | - | - |
| MINERAL | 566.525 | 1.017.882 | - | 16.434 | 44.619 | 6.420.947 | 5.395.740 | 153.762 |
| Primary Dryland Fc | 7.654 | 1.008.801 | - | 1 | 5.550 | 6.246.289 | 5.296.195 | 76.932 |
| Primary Mangrove | 558.871 | 9.081 | - | 16.433 | 39.069 | 174.658 | 99.544 | 76.830 |
| Primary Swamp Fo | - | - | - | - | - | - | - | - |
| Grand Total | 177.680.861 | 37.023.120 | 5.920.802 | 20.395.198 | 9.840.253 | 85.989.932 | 78.664.647 | 42.743.041 |
| Annual Rate | 59.226.954 | 18.511.560 | 5.920.802 | 20.395.198 | 9.840.253 | 85.989.932 | 78.664.647 | 42.743.041 |

Annex 2. Emissions From Peat Decomposition

Emissions from peat decomposition have been reported in technical annex BUR until 2017. Explanation of the calculation has also been stated in the technical annex of the BUR. The following article is only to clarify the calculation process to obtain the achievement figures in 2017.

Peat decomposition: Changing process of peat form as a result of a decline in water levels caused by deforestation and degradation activities, and land utilization.

Inherited emissions: Emission of peat decomposition will continuously occur after peatland is drained due to peat forest land conversions or land utilizations. The emissions will only stop when the peatland is completely decomposed or completely rewetted. Thus, emissions are inherited from one to another after the initial disturbance and the total emission from peat decomposition is the accumulation of peat emissions from 1990 onwards.

Emission factor for peat decomposition emission calculation: The emission factors used in the calculation are derived from the document “2013 supplement to the IPCC 2006 Guidelines for National GHG Inventory: Wetlands (2014)”.

These emission factors are used with the assumption that all utilized areas are drained. For instance, if there is a transition from primary swamp forest to secondary swamp forest, we will use the mean emission factor of the two land cover types, $(0+19)/2 = 9.5 \text{ t CO}_2 \text{ ha}^{-1}\text{yr}^{-1}$. Because it was assumed that the transition occurs gradually within the transition period, rather than abruptly in the first or the last year of the period.

There are activities needed to be seriously and continuously done for reducing the emission from peat decomposition. Those mitigation action include peat land rewetting, establishing water management systems for peat land, reducing deforestation and degradation and preventing fires on peat land.

Calculation of emissions from peat decomposition in particular year at the time of deforestation and forest degradation used the same basis as the one used in calculation of emissions from deforestation and forest degradation with the inclusion of inherited emission. As mentioned above, this is because once deforestation and forest degradation occurred in peat forests, there would be emissions from the loss of ABG at the time of conversion as described above, and additional subsequent emissions from peat decomposition at the time of deforestation and forest degradation. In addition, the deforested and degraded peat forests will release further CO₂ emissions in the following years, known as inherited emissions from peat decomposition. The emission from peat decomposition is calculated using Equation follows :

$$PDE_{ijt} = A_{ijt} \cdot EF_j$$

Where: PDE_{ijt} is Peat Decomposition Emission (PDE), i.e. CO₂ emission (tCO₂ yr⁻¹) from peat decomposition occurring in peat forest area- i that changed into land-cover type- j within time period- t ; A_{ijt} is area- i of peat forest that changed into land-cover type- j within time period- t ; EF_j is the emission factor from peat decomposition of peat forest that changed into land-cover class- j (tCO₂ ha⁻¹ yr⁻¹).

Consistent with deforestation and forest degradation activities, the emission from peat decomposition was calculated from 2013 to 2017. The base calculation for peatland emission is the area located on forested peatland in 1990. The emission baseline of peat decomposition for FREL was estimated using a linear equation approach. This estimate will be improved gradually through a stepwise process to produce a more accurate estimate for future implementation.

The decomposition process in organic soil will produce significant carbon emissions when organic soils are drained. The soils will be exposed to the aerobic condition, being oxidised and emit CO₂. In another hand, when forested peatland being converted to other land uses, the organic soils will continuously decompose for years. These emissions are inherited for years after the initial disturbance. Therefore, emissions from peat decomposition will always increase with an additional peatland being deforested. Regarding consistency, the data, methodologies, and procedures used for calculating the results presented in this report are similar to those used when establishing the FREL.

For example, in the land cover transition matrix of peatlands in the 2012-2013 period, the change of primary swamp forest (PSF) to swamp shrubs (SSr) was 3,379 ha (see Table Annex 2.1 at column 5, line 10), which was considered as the activity data. The emission factor used for this land cover transition (Table annex 2.2 at column 5, line 10), was the mean of emissions factor of the two land cover types, in this case (0+19)/2 or equals to 9.5 tCO₂/year. Thus, the emission from the peat decomposition of this deforestation was 3,379 × 9.5 equals to 32,102 ton CO₂ (see table annex 2.3. at column 5, line 10). In the following years, the emission of peat decomposition from the swamp shrubs continues as inherited emission at a rate of 19 ton CO₂/year. This rate will change if the shrubs are converted to other land use that has different emission factor.

Table Annex 2.1. Land cover transition matrix of peatlands in 2012-2013 period (in hectares)

| LC | 2012 | | | | | | | | | | | | | | | | | | | | | | | Grand Total |
|-------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|-----------|--------|---------|---------|--------|--------|---------|--------|-------|-------|---------|----|---------|--------|-------------|
| | PF | SF | PMF | SMF | PSF | SSF | TP | Sr | EP | SSr | AUA | MxUA | Rc | Sv | Po | Sw | Se | Ai | Tr | Br | Mn | WB | Ot | |
| 2013 | PF | 372,446 | | | 1 | | 16 | | | | | | | | | | | | | | | | | 372,446 |
| | SF | 4,573 | 292,000 | | | | 19 | | | | | | | | | | | | | | | | | 296,573 |
| | PMF | | | 232,928 | | | | | | | | | | | | | | | | | | | | 232,928 |
| | SMF | | | 3,847 | 89,838 | | | | | | | | | | | | | | | | | | | 93,700 |
| | PSF | | 2 | | | 2,124 | 918 | | | | | | | | | | | | | | | | | 2,124,918 |
| | SSF | | 1,145 | | 37 | 10,206 | 3,368 | 605 | 755 | 329 | 115 | 10,881 | | 224 | | | | | | | | 429 | | 3,392,605 |
| | TP | | 31 | | 50 | 585 | 27,950 | 517,985 | 4,548 | 1,420 | 19,059 | 115 | 513 | | | | | | | | | 46,389 | | 618,605 |
| | Sr | | 1,121 | | | | 1,068 | | 106,438 | | | | | 13 | | | | | | | | | | 108,560 |
| | EP | | 105 | | 10 | 15 | 19,188 | 2,226 | 2,092 | 992,893 | 42,555 | 89 | 35 | 6,467 | | | | | | | | 26,090 | | 1,091,000 |
| | SSr | | 342 | 22 | 137 | 3,379 | 57,595 | 275 | 205 | 515 | 1,791 | 213 | | | | 5,131 | | | | | | 97 | | 1,858,000 |
| | AUA | | 8,890 | | | | 1,186 | | | 598 | 1,844 | 87,988 | | | | | | | | | | 238 | | 100,776 |
| | MxUA | | 2,103 | | | | 490 | | 55,956 | | 4,378 | 2,747 | 120,391 | | | | | | | | | | | 186,147 |
| | Rc | | | | | | | | | | 33 | | | 51,552 | | | | | | | | | | 51,552 |
| | Sv | | | | | | | | | | | | | | 31,703 | | | | | | | | | 31,703 |
| | Po | | | | | | | | | | | | | | | 1,535 | | | | | | | | 1,535 |
| | Sw | | | | | | | | | | | | | | | | 95,234 | | | | | | | 95,234 |
| | Se | | | | | | | | | | | | | | | | | 5,014 | | | | | | 5,014 |
| | Ai | | | | | | | | | | | | | | | | | | 72 | | | | | 72 |
| | Tr | | | | | | | | | | | | | | | | | | | 669 | | | | 669 |
| | Br | | 959 | | 33 | 6,104 | 93,206 | 28,124 | 1,077 | 4,153 | 11,531 | 5 | 86 | 109 | | | | | | | | 320,660 | | 466,000 |
| Mn | | 28 | | | | 554 | | | | 3 | | | | | | | | | | 1,823 | | | 2,400 | |
| WB | | | | | | | | | | | | | | | | | | | | | | 824 | 824 | |
| Ot | | | | | | | | | | | | | | | | | | | | | | | - | |
| Grand Total | 377,019 | 306,728 | 363,837 | 90,102 | 145,203 | 569,878 | 549,366 | 170,447 | 999,694 | 1,881,538 | 90,983 | 121,267 | 58,128 | 31,703 | 55,100 | 365,014 | 72 | 669 | 1,823 | 393,908 | 24 | - | 11,133 | |

Table Annex 2.2. Matrix of emission factors for peat decomposition (in tCO₂/ha)

| LC | | T1 | | | | | | | | | | | | | | | | | | | | | | |
|----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|
| | | PF | SF | PMF | SMF | PSF | SSF | TP | Sr | EP | SSr | AUA | MxUA | Rc | Sv | Po | Sw | Se | Ai | Tr | Br | Mn | WB | Ot |
| TO | PF | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | SF | 9.5 | 19.0 | 9.5 | 19.0 | 9.5 | 9.5 | 46.0 | 19.0 | 29.5 | 19.0 | 35.0 | 35.0 | 27.0 | 27.0 | 9.5 | 9.5 | 27.0 | 9.5 | 35.0 | 35.0 | 9.5 | 9.5 | |
| | PMF | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | SMF | 9.5 | 19.0 | 9.5 | 19.0 | 9.5 | 9.5 | 46.0 | 19.0 | 29.5 | 19.0 | 35.0 | 35.0 | 27.0 | 27.0 | 9.5 | 9.5 | 27.0 | 9.5 | 35.0 | 35.0 | 9.5 | 9.5 | |
| | PSF | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | SSF | 9.5 | 19.0 | 9.5 | 19.0 | 9.5 | 19.0 | 46.0 | 19.0 | 29.5 | 19.0 | 35.0 | 35.0 | 27.0 | 27.0 | 9.5 | 9.5 | 27.0 | 9.5 | 35.0 | 35.0 | 9.5 | 9.5 | |
| | TP | 36.5 | 46.0 | 36.5 | 46.0 | 36.5 | 73.0 | 46.0 | 56.5 | 46.0 | 62.0 | 62.0 | 54.0 | 54.0 | 36.5 | 36.5 | 54.0 | 36.5 | 62.0 | 62.0 | 62.0 | 36.5 | 36.5 | |
| | Sr | 9.5 | 19.0 | 9.5 | 19.0 | 9.5 | 19.0 | 46.0 | 19.0 | 29.5 | 19.0 | 35.0 | 35.0 | 27.0 | 27.0 | 9.5 | 9.5 | 27.0 | 9.5 | 35.0 | 35.0 | 9.5 | 9.5 | |
| | EP | 20.0 | 29.5 | 20.0 | 29.5 | 20.0 | 29.5 | 56.5 | 29.5 | 40.0 | 29.5 | 45.5 | 45.5 | 37.5 | 37.5 | 20.0 | 20.0 | 37.5 | 20.0 | 45.5 | 45.5 | 20.0 | 20.0 | |
| | SSr | 9.5 | 19.0 | 9.5 | 19.0 | 9.5 | 19.0 | 46.0 | 19.0 | 29.5 | 19.0 | 35.0 | 35.0 | 27.0 | 27.0 | 9.5 | 9.5 | 27.0 | 9.5 | 35.0 | 35.0 | 9.5 | 9.5 | |
| | AUA | 25.5 | 35.0 | 25.5 | 35.0 | 25.5 | 35.0 | 62.0 | 35.0 | 45.5 | 35.0 | 51.0 | 51.0 | 43.0 | 43.0 | 25.5 | 25.5 | 43.0 | 25.5 | 51.0 | 51.0 | 25.5 | 25.5 | |
| | MxUA | 25.5 | 35.0 | 25.5 | 35.0 | 25.5 | 35.0 | 62.0 | 35.0 | 45.5 | 35.0 | 51.0 | 51.0 | 43.0 | 43.0 | 25.5 | 25.5 | 43.0 | 25.5 | 51.0 | 51.0 | 25.5 | 25.5 | |
| | Rc | 17.5 | 27.0 | 17.5 | 27.0 | 17.5 | 27.0 | 54.0 | 27.0 | 37.5 | 27.0 | 43.0 | 43.0 | 35.0 | 35.0 | 17.5 | 17.5 | 35.0 | 17.5 | 43.0 | 43.0 | 17.5 | 17.5 | |
| | Sv | 17.5 | 27.0 | 17.5 | 27.0 | 17.5 | 27.0 | 54.0 | 27.0 | 37.5 | 27.0 | 43.0 | 43.0 | 35.0 | 35.0 | 17.5 | 17.5 | 35.0 | 17.5 | 43.0 | 43.0 | 17.5 | 17.5 | |
| | Po | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | Sw | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | Se | 17.5 | 27.0 | 17.5 | 27.0 | 17.5 | 27.0 | 54.0 | 27.0 | 37.5 | 27.0 | 43.0 | 43.0 | 35.0 | 35.0 | 17.5 | 17.5 | 35.0 | 17.5 | 43.0 | 43.0 | 17.5 | 17.5 | |
| | Ai | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | Tr | 25.5 | 35.0 | 25.5 | 35.0 | 25.5 | 35.0 | 62.0 | 35.0 | 45.5 | 35.0 | 51.0 | 51.0 | 43.0 | 43.0 | 25.5 | 25.5 | 43.0 | 25.5 | 51.0 | 51.0 | 25.5 | 25.5 | |
| | Br | 25.5 | 35.0 | 25.5 | 35.0 | 25.5 | 35.0 | 62.0 | 35.0 | 45.5 | 35.0 | 51.0 | 51.0 | 43.0 | 43.0 | 25.5 | 25.5 | 43.0 | 25.5 | 51.0 | 51.0 | 25.5 | 25.5 | |
| | Mn | 25.5 | 35.0 | 25.5 | 35.0 | 25.5 | 35.0 | 62.0 | 35.0 | 45.5 | 35.0 | 51.0 | 51.0 | 43.0 | 43.0 | 25.5 | 25.5 | 43.0 | 25.5 | 51.0 | 51.0 | 25.5 | 25.5 | |
| | WB | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |
| | Ot | - | 9.5 | - | 9.5 | - | 9.5 | 36.5 | 9.5 | 20.0 | 9.5 | 25.5 | 25.5 | 17.5 | 17.5 | - | - | 17.5 | - | 25.5 | 25.5 | - | - | |

Table Annex 2.3. Matrix of CO₂ emissions from peat decomposition (in tCO₂e)

| LC | | 2012 | | | | | | | | | | | | | | | | | | | | Grand Tot | | | |
|-----------|--------|-----------|-----------|---------|---------|------------|------------|------------|-----------|-----------|-------------|---------|-----------|-----------|-----------|--------|---------|---------|--------|--------|------------|------------|-----------|----------|----------|
| | | PF | SF | PMF | SMF | PSF | SSF | TP | Sr | EP | SSr | AUA | MxUA | Rc | Sv | Po | Sw | Se | Al | Tr | Br | | Mn | WB | Ot |
| 2013 | PF | - | - | - | 5 | - | 150 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 150 |
| | SF | 43,442 | 5,548,000 | - | - | - | 358 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 5,591,8 |
| | PMF | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| | SMF | - | - | 36,941 | 706,913 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1,743,8 |
| | PSF | - | 16 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 16 |
| | SSF | - | 21,730 | - | 703 | 96,992 | 64,003,498 | 34,747 | 6,260 | 3,388 | 206,744 | - | 7,832 | - | - | - | - | - | - | - | - | - | 15,001 | - | 64,396,8 |
| | TP | - | 1,434 | - | 2,300 | 21,343 | - | 37,812,876 | 209,100 | 80,255 | 876,716 | 7,108 | 32,137 | - | - | - | - | - | - | - | - | - | 2,876,095 | - | 41,919,8 |
| | Sr | - | 21,306 | - | - | - | 20,297 | - | 2,022,327 | - | - | - | 447 | - | - | - | - | - | - | - | - | - | - | - | 2,064,3 |
| | EP | - | 3,094 | - | 308 | 308 | 566,045 | 125,751 | 61,716 | 39,715 | 301,255,384 | 4,051 | 1,592 | 242,497 | - | - | - | - | - | - | - | - | 1,187,080 | - | 43,163,3 |
| | SSr | - | 6,506 | 212 | 2,611 | 32,102 | 1,094,303 | 12,692 | 3,920 | 15,182 | 34,033,441 | - | - | - | - | - | 48,743 | - | - | - | - | - | 3,387 | - | 35,252,7 |
| | AUA | - | 311,140 | - | - | - | 41,543 | - | 27,222 | 65,997 | 4,487,368 | - | - | - | - | - | - | - | - | - | - | - | 12,127 | - | 4,945,3 |
| | MxUA | - | 73,599 | - | - | - | 17,119 | - | 1,958,469 | - | 153,225 | 142,156 | 139,445 | - | - | - | - | - | - | - | - | - | - | - | 8,484,5 |
| | Rc | - | - | - | - | - | - | - | - | - | 897 | - | - | 1,804,327 | - | - | - | - | - | - | - | - | - | - | 1,805,2 |
| | Sv | - | - | - | - | - | - | - | - | - | - | - | - | - | 1,109,621 | - | - | - | - | - | - | - | - | - | 1,109,6 |
| | Po | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| | Sw | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| | Se | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 175,494 | - | - | - | - | - | - | 175,4 |
| | Al | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Tr | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 34,130 | - | - | - | 34,13 | |
| Br | - | 33,552 | - | 1,147 | 155,660 | 3,262,111 | 743,714 | 37,684 | 188,941 | 403,577 | 267 | 4,378 | 4,688 | - | - | - | - | - | - | - | - | 16,353,669 | - | 22,189,9 | |
| Mn | - | 991 | - | - | - | 19,386 | - | - | - | 97 | - | - | - | - | - | - | - | - | - | - | - | 92,961 | - | 113,4 | |
| WB | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Ot | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Grand Tot | 43,442 | 6,021,388 | 37,131 | 713,987 | 306,336 | 69,024,939 | 729,194 | 299,560 | 0,030,718 | 6,995,438 | 640,516 | 186,322 | 2,051,513 | 1,109,621 | - | 48,743 | 175,494 | - | 34,130 | 92,961 | 20,447,359 | - | - | 232,990 | |

Historical emission from peat decomposition

The emissions from peat decomposition is progressive, due to inherited emissions from previous degraded peatlands. The emissions from peat decomposition will never decrease unless the degraded peatlands are changed into forests, which is unlikely to happen in this period of assessment. In the first FREL document, we developed linear equations from regression analysis using annual peat emissions from historical data. The emissions from peat decomposition were estimated based on the land cover maps. In some years, instead of yearly land cover map, we only have multi-years land cover maps, i.e. 3-yearly (2006 – 2009, 2-yearly (2009-2011) and 1-yearly (2011-2016). We generated annual emission from the average values of the mapping period. Each year has an estimated emission value to be regressed against year.

For the construction of reference emission level 2017-2020, consistent with method in first FREL document used linear projection with equation $y = 6.706.744,03x - 13.266.946.368,06$ $R^2 = 0,97$ The reference period 2006/2007 – 2015/2016 (see table annex 2.4 and figure annex 2.1)

Table annex 2.4. Emission from peat decomposition

| Year | Peat Decomposition | Actual Emission |
|-----------|--------------------|---|
| 2006-2007 | 200.067.598 | Historical |
| 2007-2008 | 200.067.598 | |
| 2008-2009 | 200.067.598 | |
| 2009-2010 | 215.742.080 | |
| 2010-2011 | 215.742.080 | |
| 2011-2012 | 226.109.789 | |
| 2012-2013 | 234.152.020 | |
| 2013-2014 | 240.799.350 | |
| 2014-2015 | 248.530.578 | |
| 2015-2016 | 255.413.778 | |
| 2016-2017 | 260.556.280 | 256.694.322 |
| 2017-2018 | 267.263.024 | Projection $y = 6.706.744,03x - 13.266.946.368,06$ $R^2 = 0,97$ |
| 2018-2019 | 273.969.768 | |
| 2019-2020 | 280.676.512 | |
| | | |




Figure Annex 2.1. The emissions from peat decomposition

Emission Reduction from peat decomposition in 2017

Peat decomposition emissions in 2017 (actual emission 256.694.322 tCO₂) , when compared to the historical emissions projections in the reference emission level 2006-2016 (projected in 2017 is 260.556.280 tCO₂) , then the **emission reduction in 2017 is 3.861.958 tCO₂**

Annex 3. Emissions From Peat Fires

Emissions from peat fires have not been included in Indonesia's first FREL calculation. Peat fire emission data is presented in the annex of the first FREL document. However, in the protocol MRV document, emissions from peat fires were also reported. Until now, no reference has been made for peat fires. Thus, the reference to peat fires will be made using peat fire data from 2006-2016. The method and data used for this report are consistent with the annex first FREL document regarding peat fire.

According to the IPCC Supplement for Wetland (IPCC, 2014), emissions from organic soil fires are calculated with the following formula:

$$E_{\text{peat}} = A \times MB \times CF \times G_{\text{ef}}$$

Where, E_{peat} is emission from peat fires, A is burned peat area, MB is mass of fuel available for combustion, CF is combustion factor (default factor = 1.0) and G_{ef} is emissions factor.

Fire activity data from 2006-2014 used a method developed by the Mitsubishi Research Institute (MRI) with a 1 x 1 km grid approach.

Tier 1 estimation of peat fire emission requires data on burn scar area. The currently available methods for determining burned scar area are based on low resolution MODIS images or hotspots analysis (MRI, 2013). However, the MODIS collection 5 of burned areas (MCD45A1) data had no observation over SE Asia regions, especially for major Islands of Indonesia.

The following is the method adapted from MRI (2013) to generate burn scar map in peatland based on hotspot analysis. The method was developed from a REDD+ demonstration activity project in Central Kalimantan. First, hotspots data are compiled annually from the baseline years (e.g. 1990, 1991, 1992, 1993, etc.). To improve certainty, only hotspots with confidence level of more than 80% are selected. As MODIS hotspots are not available for the period before 2000, NOAA hotspot might be used for to fill the gap. However, comparability and accuracy of NOAA hotspots need to be assessed, as they do not have the information on the confidence level. Second, a raster map with 1x1 km grid (pixel size) is generated and overlaid on top of the hotspot data. Pixels without hotspots are considered as not burned and excluded from the activity data. Each 1km x1 km pixel with at least one hotspot is considered as burned but with the assumption that the burned area is 75% of the pixel area (7,500 ha). This rule applies for each pixel regardless the number of hotspots within a particular pixel (Figure Annex 4.1). Then, these burned areas were overlaid with the peat land map (produced by MoA) to estimate the burned peat land for each year.




Figure Annex 3.1. Methodology to derive burned area (activity data)

While for 2015 - 2017 fire activity data uses visual interpretation methods, making it more accurate. Based on indications of hotspots with more than 80% confidence level, point density analysis was made. This is to make the initial polygon area burn. Furthermore Landsat with coverage dates after the fire (max. 14 days afterwards) is used as a reference to digitize actual burn scar. As an illustration see the picture ...




Figure Annex 3.2. Delineation of burn scar

Mass of fuel available for combustion

Mass of fuel available for combustion, MB, is estimated from multiplication of mean depth of burned peat (D) and bulk density (BD), assuming average peat depth burned by fire is 0.33 m (Ballhorn *et al.*, 2009) and bulk density is 0.153 ton/m³ (Mulyani *et al.*, 2012). Resulted mass available for combustion is 0.05049 ton/m² or 504.9 ton/ha.

Emission factor

CO₂ emission factor (kg CO₂/kg) can be indirectly estimated from organic carbon content (kg C/kg of weight), which is equal to:

$$E_{CO_2} = C_{org} \times 3.67$$

C_{org} can be estimated by the following equation:

$$C_{org} = \frac{(1 - \frac{M_{ash}}{M_s})}{1.724} \times 3.67$$

Where M_s is mass of soil solids, which is equal to accumulation mass of ash (M_{ash}) and mass of organic matters. Ratio of M_{ash} and M_s is 14.04%, which is the mean ash contents of three peat types; namely, Sapric (4.98%), Hemic (21.28%) and Fibric (15.85%) (see Mulyani *et al.*, 2012).

Adjustment factor of 1/1.724 is used to convert organic matter estimate to organic carbon content. Estimated C_{org} is 49.86% (or kg/kg), which is equal to 498.6 C g/kg dry matter burnt.

If the value is converted to CO₂e, the value would be $C_{org} \times 3.67 = 1,828.2$ CO₂ g/kg dry matter burnt or 1,828.2 CO₂ kg/ton. Assuming of 1 ha peat burning, CO₂ emissions released to the atmosphere is:

$$\begin{aligned} E_{CO_2} &= A \times Y \times E_{CO_2} \\ &= 1 \text{ ha} \times 504.9 \text{ t/ha} \times 1,828.2 \text{ kg/t} \\ &= 923,058.18 \text{ kg/ha} \\ &= 923.1 \text{ tCO}_2\text{e/ha} \end{aligned}$$

This result is used as emission factor of burned peat. Emissions from peatlands that suffer more than one fire event are assumed to be reduced by half compared to that of the first burning, e.g. the first burning of 1 ha peat emits 923.1 tCO₂ (UKP4 and UNORCID, 2013), while the subsequent burning of exactly the same area will release 462 tCO₂. The third burning of the same area will release again a lower amount of emissions than the second burning but further research is necessary to determine the amount of reduction. The above assumption is from a manuscript that resulted from Peat Emission Workshop held by UKP4 and UNORCID (6 November 2013) in Jakarta.

Historical emission from peat fire

For this report, historical emissions from peat fire have been calculated for the period 2006-2016.

It was found that the annual estimated burned peat areas varied greatly from 2006 to 2016 (Figure Annex 3.3). The highest rate occurred in 2015 accounting for 869,754 ha of burned peatland, while the lowest rate occurred in 2010 accounting for 55,664 ha of burned peat area. Using this historical data set, the average value used as activity data for proposed REL from burned peat accounts for 269,686 ha.




Figure Annex 3.3. Estimated burned peat area

The results of the calculation of emissions from burned peat are shown in Figure Annex 3.4. Average emission from peat fire from 2006 – 2016 is 248,947,149 tCO₂e yr⁻¹. The derivation of the burned areas has not been verified using ground truthing or high-resolution satellite data. Therefore, the uncertainty level cannot be estimated.




Figure Annex 3.4. Estimated historical emission from burned peat

Emission reduction from peat fire in 2017

Emissions from peat fires in 2017 have decreased dramatically, this was due to massive law enforcement and moratorium policies for new permits on peat and primary forests. The fire incident in peat in 2017 was 13,555 hectares. If it is converted to emissions, it will amount to 12,512,621 tCO₂. When compared to the reference emission level for peat fire (248,947,149 tCO₂), the **emission reduction in 2017 is 236,434,529 tCO₂**.

Constraints in measuring emissions from peat fires

Some critical issues on the accuracy of the burn scar lies in the assumptions used to estimate the size and intensity of the fires. Hotspots are just an indication of active fire existence through thermal differentiation with neighboring pixels. Thus, false detection is possible as a thermal anomaly can originate from other heat sources than fires. Selection of hotspot with high confidence level can reduce such error. However, smoke coverage is very common during fire season, which reduces the sensor's capability to detect fires covered by smokes. This can result in underestimation of the burned areas. In contrast, assuming that the burned area is 75% for each pixel with hotspot might lead to a severe overestimate of the burned area, especially in the border area between burned and unburned.

A further challenge lies in determining the peat depth consumed by fires. Relationship analyses between hotspot parameters (fire intensity, frequency etc.) with burned peat depth need to be carried out to better estimate the burned peat depth of the burned peatland and thus estimate the actual emissions from peat fires. Ballhorn *et al.* (2009) used airborne LIDAR for estimating burned peat depth with accuracy of less than 20 cm. Konecny *et al.* (2016) found that carbon loss varies significantly for recurrent fires in drained tropical peatlands. According to their research the relative burned area depth decreases over the first four fire events and is then constant for further successive fires. They estimate values for the dry mass of peat fuel consumed to be only 58–7% of the current IPCC Tier 1 default value for all fires. This indicates that accurate estimation of emissions from peat fires should also consider the frequency of fires in an area and employ accordingly adjusted emission factors.

Improvement of Peat Fire Data used the data burn area from Ministry of Environment and Forestry with the a new approach for estimating the burned area. This improved method has been applied for estimation of the burn scar, i.e. by combining the Landsat image (quick look original with composite band 645) with the hotspot data and verified with observed burnt area data on the ground. That is able to delineate the burn area. This new approach might be adopted in the future as this approach will have higher certainty.

With above conditions and high level of uncertainties of all involved parameters (hotspot detection, size of burned area estimation, fire frequency, burned peat depth, mass of fuel available for combustion), this FREL document did not include emissions from peat fires. Advancing technology in remote sensing to improve burned scar and peat depth mapping will increase the accuracy of peat fire emission calculation which can then be included as improvement in a future FREL.

Annexes References

- Ballhorn, U., Siegert, F., Mason, M. and Limin, S. 2009 Derivation of burn scar depths and estimation of carbon emissions with LIDAR in Indonesian peatlands. *Proceedings of the National Academy of Sciences of the United States of America*, 106 (50), 21213-21218.
- Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. 2014. *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands* (p. 354). Intergovernmental Panel on Climate Change: (IPPC), Switzerland.
- Konecny, K., Ballhorn, U., Navratil, P., Jubanski, J., Page, S.E., Tansey, K., Hooijer, A., Vernimmen, R. and Siegert, F. 2016. *Variable carbon losses from recurrent fires in drained tropical peatlands*. *Global Change Biology* 22, pp. 1469–1480.
- MRI 2013. *Green House Gas Reduction Project through Forest Conservation in Peat land in Central Kalimantan*: Mitsubishi Research Institute, Inc.
- Mulyani, et al. 2012. *Basis Data Karakteristik Tanah Gambut Di Indonesia*. Prosiding Seminar Nasional Pengelolaan Lahan Gambut Berkelanjutan 4 Mei 2012: From <http://balittanah.litbang.deptan.go.id>